

Acute toxicity effect of glyphosate on survival rate of common carp, *Cyprinus carpio*

Mohammad Forouhar Vajargah^{1*}, Ahmad Mohamadi Yalsuyi², Masoud Sattari^{1,3}, Aliakbar Hedayati⁴

¹Department of Fisheries, School of Natural Resources, University of Guilan Sowmehsara, Sowmehsara, Iran

²Department of Aquaculture, School of Fisheries and Environment, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

³Department of Marine Sciences, Caspian Sea Basin Research Center, University of Guilan, Rasht, Iran

⁴Department of Aquatic Production and Exploitation, School of Fisheries and Environment, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

Abstract

Background: Herbicides are usually used to control weeds and some of them like glyphosate are non-selective herbicides. Aquatic environments are usually the last destinations of agricultural pesticides, which disrupt the metabolic processes of organisms. Thus, the aim of this study was to evaluate acute toxicity of glyphosate on the survival rate of common carp, *Cyprinus carpio*.

Methods: A total of 135 common carp averaged 7 ± 0.8 g in weight were exposed to 0, 25, 50, 75, 100, 125, 150, 175 and 200 mL.L⁻¹ glyphosate (15 fish in each treatment, with triplicates) for 96 hours. The aquariums capacity was 98 L in volume (80 × 35 × 35 cm) and physicochemical parameters were the same for all groups (pH 7.4-8, temperature = $26 \pm 1^\circ\text{C}$, DO = 7 mg.L⁻¹ and total hardness of 190 mg CaCO₃). LC₁₀, LC₂₀, LC₃₀, LC₄₀, LC₅₀, LC₆₀, LC₇₀, LC₈₀, LC₉₀ and LC₉₅ of glyphosate were calculated at 24, 48, 72 and 96 hours after adding glyphosate using probit test.

Results: Mortality was observed in all treatments which exposed to higher than 50 mL.L⁻¹ after 96 hours. The results showed that 96-hour LC₅₀ of glyphosate for common carp was 92.71 mL.L⁻¹. The fish exposed to different concentrations of glyphosate showed clinical signs such as increased mucus secretion, skin darkening and death with mouth open.

Conclusion: Glyphosate disrupts the synthesis of amino acids in plants by inhibiting enzymatic activity of 5-enolpyruvylshikimate-3-phosphate synthase (EPSP). This enzyme is absent in animals. However, glyphosate is toxic for common carp.

Keywords: Animals, Herbicides, Glyphosate, Carps, Excitatory postsynaptic potentials

Citation: Forouhar Vajargah M, Mohamadi Yalsuyi A, Sattari M, Hedayati A. Acute toxicity effect of glyphosate on survival rate of common carp, *Cyprinus carpio*. Environmental Health Engineering and Management Journal 2018; 5(2): 61–66. doi: 10.15171/EHEM.2018.09.

Article History:

Received: 29 November 2017

Accepted: 3 February 2018

ePublished: 28 February 2018

*Correspondence to:

Mohammad Forouhar Vajargah

Email:

mohammad.forouhar@yahoo.com

Introduction

In recent years, the use of agricultural pesticides has been increased significantly (1). Various types of agricultural pesticides such as herbicides, insecticides and fungicides are commonly used globally (2); these pesticides usually spread in the aquatic environments through various ways such as rainfall and soil erosion (3). In other words, aquatic ecosystems are the ultimate destinations of agricultural pesticides (4). Aquatic organisms contact with their environment through physiological surfaces and pesticides can penetrate through the skin, gills and mouth into their body, and affect liver, kidneys, gonads and other organs (5). Contamination and accumulation of pesticides in the aquatic environment can lead to food

poisoning and mortality in humans (1,2).

Herbicides are commonly used in farms to control weeds (6). In the meantime, glyphosate (glyphosate 41% SL) is one of the most widely used herbicides in many countries, especially developing countries (1,7). The use of glyphosate had the fastest growth rate compared to other herbicides (8). Two-thirds of the total volume of glyphosate has been sprayed in the recent decades (9). Glyphosate inhibits the 5-enolpyruvylshikimate-3-phosphate synthase (EPSP) activity in plants and impairs the synthesis of the aromatic amino acids (10). EPSP is found only in plants and some bacteria (11).

Glyphosate is highly soluble in water (10500 mL.L⁻¹) and its half-life, depending on the environmental conditions,



is between 3.5 and 70 days (12). However, Souza et al (13) pointed out that the half-life of glyphosate in freshwater was between 28 and 87 days. It was thought that since EPSP enzyme was not found in animals, so glyphosate is not toxic to them. However, the results of previous studies showed that it was toxic to fish species (10,14). Loro et al (15) reported that glyphosate-based herbicide was toxic to *Rhamdia quelen* and *Leporinus obtusidens*. Sandun et al (16) reported that 96-hour LC_{50} of glyphosate to *Oreochromis niloticus* was 16.7 mg.L^{-1} ; Da Cruz et al (17) also pointed out that glyphosate-based herbicide was toxic to fish as an aquatic organism.

Fish, as aquatic organisms, can be used as biomarkers in toxicology and biological studies (18). So fish often have been used as experimental biological models for exhibiting environmental impacts of toxins (19). Toxicological studies on lethal concentration of toxins are very important (20), and one of the methods is to determine 96-hour LC_{50} of pesticides on fish (21,22). Toxicological studies like those determining 96-hour LC_{50} of toxins on fish can provide better management of water resources as a method for detection of biological effects of pollutants (23). Finally, these studies can also be useful in determining the limits of toxins in nature and human food safety (24).

Common Carp, *Cyprinus carpio* Linnaeus, 1758, as a commercially- important fish species in aquaculture (25) is found in most temperate regions of the world (26). It is omnivorous (27) and often searches bottom of the pond for getting food (28). Feeding habits, high growth rate, reproduction in captivity, adaptation to commercial diets and density are reasons for the common carp popularity in aquaculture industries (29). However, human activities and environmental pollutants can disrupt the life cycle of common carp (30). The use of herbicides such as glyphosate in addition to the destruction of macrophytes may affect non-target organisms such as common carp through destruction of habitats and food (31,32). For example Zebrafish, *Danio rerio*, needs a substrate full of plants for spawning and reproduction (33), so the pollutants may have negative effects on breeding of this species through the destruction of non-target aquatic plants.

There is limited information about the toxicity of glyphosate. It is also true for the maintenance of carp in ponds and reservoirs as well as its long half-life in the aquatic environments. So, the aim of the present study was to assess acute toxicity of glyphosate on survival rate of common carp, *C. carpio* as an animal model.

Methods

The experiment was performed according to Yalsuyi and Vajargah (2) and Montajami et al (34).

Sample preparation

200 common carp averaged $7 \pm 0.8 \text{ g}$ in weight were prepared from some farms in Guilan province and

transferred to the laboratory in Faculty of Natural Resources, University of Guilan, Guilan province, Iran, and divided into four 250-L tanks (50 fish in each tank). Fish were maintained in these tanks for two weeks in order to acclimate with laboratory conditions. Fish were fed a commercial diet (produced by Faradaneh Co. Tehran, Iran) at 3% of body weight 3 times a day. Physicochemical parameters of water in all tanks were the same (pH 7.4-8, temperature = $26 \pm 1^\circ\text{C}$, DO = 7 mg.L^{-1} and total hardness = 190 mg CaCO_3).

Toxicity tests

After adaptation, 135 fish were selected randomly and divided into 9 treatment groups (0, 25, 50, 75, 100, 125, 150, 175 and 200 mL.L^{-1} glyphosate) in triplicates into 27 aquariums ($80 \times 35 \times 35 \text{ cm}$, capacity 85 l). Fish were exposed to different concentrations of glyphosate (Glyphosate Aria 41% SL) for 96 hours. Mortality rates of fish were recorded at 24, 48, 72 and 96 hours after adding glyphosate. Animals then were transferred into aquariums 16 hours before the toxicity test and feeding was stopped 24 hours before the experiment. Water physicochemical parameters were the same as the adaptation time (pH 7.4-8, temperature = $26 \pm 1^\circ\text{C}$, DO = 7 mg.L^{-1} and total hardness = 190 mg CaCO_3).

Data analysis

Toxicity of glyphosate was estimated using the method proposed by Vajargah and Hedayati (35). Lethal concentrations of glyphosate for 50% of the population at 24, 48, 72 and 96 h LC_{50} were calculated using probit test (with a 95% CI) by IBM SPSS 20. Spearman test (two-tailed test) was used to determine the correlation between different concentrations of glyphosate and fish mortality.

Results

Mortality was not observed in the adaptation period. The results of toxicity tests showed that there was a significant correlation ($P < 0.01$) between concentration of glyphosate and fish mortality. All fish died after 96 hours exposure to the concentrations higher than 150 mL.L^{-1} glyphosate, while in concentration of 200 mL.L^{-1} , all fish died after 24 hours (Table 1).

96-h LC_{50} of glyphosate for common carp was 92.711 mL.L^{-1} . LC_{10} , LC_{20} , LC_{30} , LC_{40} , LC_{50} , LC_{60} , LC_{70} , LC_{80} , LC_{90} and LC_{95} of glyphosate within 24, 48, 72 and 96 hours are demonstrated in Table 2. The fish exposed to different concentrations of glyphosate showed clinical signs such as increased mucus secretion, skin darkening, fast swimming, increased movements of operculum and death with mouth open.

Discussion

As it is estimated that the world population increases up to 9 billion persons by 2050 (36), therefore, food supply needs to be increased by productivity (37); and

Table 1. Mortality rate of common carp, *Cyprinus carpio* exposed to different concentrations of glyphosate (n=15 in each treatment)

Concentration (mg.L ⁻¹)	Number	No. of mortality			
		24 h	48 h	72 h	96 h
0	15	0	0	0	0
25	15	0	0	0	0
50	15	0	0	0	1
75	15	1	3	4	6
100	15	2	5	7	9
125	15	2	6	8	12
150	15	3	8	13	15
175	15	6	10	14	15
200	15	15	15	15	15

All fish died at concentrations higher than 150 mL.L⁻¹.

Table 2. Lethal concentration of glyphosate for common carp, *Cyprinus carpio*

Point	Concentration (mg.L ⁻¹)			
	24 h	48 h	72 h	96 h
LC ₁₀	109.420	71.438	63.006	52.731
LC ₂₀	128.898	93.337	79.563	66.455
LC ₃₀	142.943	109.130	91.501	76.352
LC ₄₀	154.944	122.624	101.702	84.808
LC ₅₀	166.161	135.236	111.237	92.711
LC ₆₀	177.378	147.848	120.771	100.615
LC ₇₀	189.379	161.342	130.972	109.071
LC ₈₀	203.425	177.135	142.911	118.967
LC ₉₀	222.903	199.036	159.467	132.691
LC ₉₅	238.988	217.122	173.40	144.025

agricultural growth requires an increase in productivity through the use of a variety of nutrients for plants and soil, improving agricultural techniques, and the use of high-yielding plant varieties and pesticides (38). Glyphosate is widely used in agriculture (9). According to the World Health Organization (WHO) (39), application of glyphosate has increased annually. It has also long half-life in aquatic environments (13) and its residues can remain stable in foods for a year or even more, even if the foods are frozen, dried or processed (40). It also damages macrophytes as well as non-target organisms such as common carp through deterioration of fish habitats and food (31). Hence, one of the main challenges concerning the widespread application of agricultural pesticides can be related to their devastating impact on non-target organisms (41).

Akinsorotan (42) reported that the 96-h LC₅₀ of glyphosate on fingerlings of *Clarias gariepinus*, as an aquatic organism, was 18.07 mg.L⁻¹. It was also found that glyphosate can reduce the survival rate of the fish, which is consistent with the results of the present study.

Micah et al (43) studied behavioral responses of *Heteroclaris* (hybrid) exposed to two types of glyphosate and found that it was toxic to *Heteroclaris* (hybrid) as an aquatic organism and its 96-hour LC₅₀ was 6.838 mg.L⁻¹.

They also reported that glyphosate had some clinical signs such as increased opercular ventilation and also caudal fin damage, which was consistent with the results of the present study.

Da Cruz et al (17) investigated sensitivity, ecotoxicity and histopathological effects on some Neotropical fish exposed to glyphosate alone and with surfactant. Their results showed that glyphosate was toxic to these species and there was significant correlation between mortality rate of fish and concentration of glyphosate ($P < 0.01$). It also reduced the survival rate of fish as non-target organism and had harmful effects on fish, which was also similar to the results of the present study.

Jofré et al (12) studied toxicity of two commercial glyphosate on survival rate of two fish species (*Danio rerio* and *Poecilia reticulata*). The results of their study indicated that both herbicides may produce potential environmental damages, which is similar to the results of the present study.

Hena Muñoz et al (44) studied acute toxicity and sub-lethal effects of the mixture glyphosate (Roundup® Active and Cosmo-Flux 411F) to anuran embryos and tadpoles of four Colombian species. The results showed that 96-hour LC₅₀ of glyphosate to embryos and tadpoles was between 2.2-3.9 µg.L⁻¹ and 1.4-2.8 µg.L⁻¹, respectively. Tadpoles of

Hypsiboas crepitans was the most sensitive and glyphosate was toxic to aquatic organisms, which is consistent with the results of the present study.

By comparing the results of the present study and Jofré et al (12), Akinsorotan (42), Micah et al (43) and Henao Muñoz et al (44), it was found that the toxicity of glyphosate depends on the size, species and physicochemical parameters of the environment.

Conclusion

According to the results, although glyphosate inhibits the enzyme EPSP and this enzyme has been observed only in plants and some bacteria, but glyphosate was toxic to *C. carpio* as a non-target organism. The results of the present study can be useful for further ecotoxicological studies. On the other hand, agricultural pesticides are associated with materials which have different combinations such as salts and a surfactant called polyethylene amine (POEA). These compounds in some cases, can increase the toxicity of pesticides. For example, Peyote[®], which is one of the glyphosate compounds, is more toxic than glyphosate. POEA is also toxic to aquatics (45). The present study can also be useful for further investigations on the toxicity of POEA to the other aquatics.

Acknowledgments

The present study was carried out by laboratory facilities and research equipment of Guilan University. The authors would like to gratitude Dr. Rasoul Ghorbani for his help and guide. We also thank all personnel of University of Guilan who helped us to carry out this laboratory examinations.

Ethical issues

Fish was used as a model of aquatic organism to study the toxicity of different pollutants. In term of ethical issues, the procedures described in this paper are in accordance with the ecological effects of test guidelines (46) published by the United States Environmental Protection Agency (US EPA).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Authors were participated in all aspects of this research work, like data collection, analysis, interpretation and manuscript approval.

References

- de Oliveira JL, Campos EV, Bakshi M, Abhilash PC, Fraceto LF. Application of nanotechnology for the encapsulation of botanical insecticides for sustainable agriculture: prospects and promises. *Biotechnol Adv* 2014; 32(8): 1550-61. doi: 10.1016/j.biotechadv.2014.10.010.
- Mohamadi Yalsuyi A, Forouhar Vajargah M. Acute toxicity of silver nanoparticles in Roach (*Rutilus rutilus*) and Goldfish (*Carassius auratus*). *J Environ Treat Tech* 2017; 5(1): 1-4.
- Larras F, Montuelle B, Bouchez A. Assessment of toxicity thresholds in aquatic environments: does benthic growth of diatoms affect their exposure and sensitivity to herbicides? *Sci Total Environ* 2013; 463-464: 469-77. doi: 10.1016/j.scitotenv.2013.06.063.
- Hedayati A, Forouhar Vajargah M, Mohamadi Yalsuyi A, Abarghouei S, Hajiahmadyan M. Acute toxicity test of pesticide abamectin on common carp (*Cyprinus carpio*). *J Coast Life Med* 2014; 2(11): 841-44. doi: 10.12980/JCLM.2.201414J44.
- Ibrahim AT, Harabawy AS. Sublethal toxicity of carbofuran on the African catfish *Clarias gariepinus*: Hormonal, enzymatic and antioxidant responses. *Ecotoxicol Environ Saf* 2014; 106: 33-9. doi: 10.1016/j.ecoenv.2014.04.032.
- Dentzman K, Gunderson R, Jussaume R. Techno-optimism as a barrier to overcoming herbicide resistance: Comparing farmer perceptions of the future potential of herbicides. *J Rural Stud* 2016; 48: 22-32. doi: 10.1016/j.jrurstud.2016.09.006.
- Poiger T, Buerge IJ, Bachli A, Muller MD, Balmer ME. Occurrence of the herbicide glyphosate and its metabolite AMPA in surface waters in Switzerland determined with on-line solid phase extraction LC-MS/MS. *Environ Sci Pollut Res Int* 2017; 24(2): 1588-96. doi: 10.1007/s11356-016-7835-2.
- Myers JP, Antoniou MN, Blumberg B, Carroll L, Colborn T, Everett LG, et al. Concerns over use of glyphosate-based herbicides and risks associated with exposures: a consensus statement. *Environ Health* 2016; 15: 19. doi: 10.1186/s12940-016-0117-0.
- Benbrook CM. Trends in glyphosate herbicide use in the United States and globally. *Environ Sci Eur* 2016; 28(1): 3. doi: 10.1186/s12302-016-0070-0.
- Tian YS, Xu J, Xing XJ, Zhao W, Fu XY, Peng RH, et al. Improved glyphosate resistance of 5-enolpyruvylshikimate-3-phosphate synthase from *Vitis vinifera* in transgenic *Arabidopsis* and rice by DNA shuffling. *Mol Breed* 2015; 35(7): 148. doi: 10.1007/s11032-015-0327-0.
- Tohge T, Watanabe M, Hoefgen R, Fernie AR. Shikimate and phenylalanine biosynthesis in the green lineage. *Front Plant Sci* 2013; 4: 62. doi: 10.3389/fpls.2013.00062.
- Jofré DM, Garcia MJ, Salcedo R, Morales M, Alvarez M, Enriz D, et al. Fish toxicity of commercial herbicides formulated with glyphosate. *J Environ Anal Toxicol* 2013; 4(1): 1-3. doi: 10.4172/2161-0525.1000199.
- Souza EL, FOLONI LL, Filho JT, Velini ED, Siono LM, Silva JR. Half-life of glyphosate on the control of water hyacinths in water tanks. *J Water Resource Prot* 2017; 9(5): 470-81. doi: 10.4236/jwarp.2017.95030.
- Bohn T, Rover CM, Semenchuk PR. *Daphnia magna* negatively affected by chronic exposure to purified Cry-toxins. *Food Chem Toxicol* 2016; 91: 130-40. doi: 10.1016/j.fct.2016.03.009.
- Loro VL, Gluszcak L, Moraes BS, Leal CAM, Menezes

- C, Murussi CR, et al. Glyphosate-based herbicide affects biochemical parameters in *Rhamdia quelen* (Quoy & Gaimard, 1824 and) *Leporinus obtusidens* (Valenciennes, 1837). *Neotrop Ichthyol* 2015; 13(1): 229-36. doi: 10.1590/1982-0224-20140082.
16. Sandun KV, Bandara N, Amarasinghe US. Effect of glyphosate-based herbicide, Roundup™ on territory deference of male *Oreochromis mossambicus* (Osteichthyes, Cichlidae) associated with mating behaviour. *Sri Lanka J Soc Sci* 2015; 20(1): 1-10. doi: 10.4038/slj.s.v20i1.7451.
 17. Da Cruz C, Carraschi SP, Shiojiri NS, Da Silva AF, Pitelli RA, Machado MR. Sensitivity, ecotoxicity and histopathological effects on neotropical fish exposed to glyphosate alone and associated to surfactant. *Journal of Environmental Chemistry and Ecotoxicology* 2016; 8(3): 25-33. doi: 10.5897/JECE2015.0362.
 18. Ullah S, Zorriehzakra MJ. Ecotoxicology: a review of pesticides induced toxicity in Fish. *Adv Anim Vet Sci* 2014; 3(1): 40-57. doi: 10.14737/journal.aavs/2015/3.1.40.57.
 19. Palanikumar L, Kumaraguru AK, Ramakritinan CM, Anand M. Toxicity, biochemical and clastogenic response of chlorpyrifos and carbendazim in milkfish *Chanos chanos*. *Int J Environ Sci Technol* 2014; 11(3): 765-74. doi: 10.1007/s13762-013-0264-6.
 20. Arome D, Chinedu E. The importance of toxicity testing. *J Pharm Biosci* 2013; 4: 146-48.
 21. Mohamadi Yalsuyi A, Forouhar Vajargah M. Recent advance on aspect of fisheries: A review. *J Coast Life Med* 2017; 5(4): 141-48. doi: 10.12980/jclm.5.2017J6-226.
 22. Forouhar Vajargah M, Hedayati A, Mohamadi Yalsuyi A, Abarghouei S, Gerami MH, Ghaffari farsani H. Acute toxicity of Butachlor to Caspian Kutum (*Rutilus frisii kutum* Kamensky, 1991). *J Environ Treat Tech* 2014; 2(4): 155-57.
 23. Forouhar Vajargah M, Hedayati A. Acute toxicity of trichlorofon on four viviparous fish: *Poecilia latipinna*, *Poecilia reticulata*, *Gambusia holbrooki* and *Xiphophorus helleri* (Cyprinodontiformes: Poeciliidae). *J Coast Life Med* 2014; 2(7): 511-14. doi: 10.12980/JCLM.2.2014J11.
 24. Khan A, Shah N, Gul A, Sahar N, Ismail A, Muhammad M, et al. Comparative study of toxicological impinge of glyphosate and atrazine (herbicide) on stress biomarkers; blood biochemical and haematological parameters of the freshwater common carp (*Cyprinus carpio*). *Polish Journal of Environmental Studies* 2016; 25(5): 1995-2001. doi: 10.15244/pjoes/62698.
 25. Shirali S, Erfani Majd N, Mesbah M, Seifi MR. Histological studies of common carp ovarian development during breeding season in Khouzestan province, Iran. *World Journal of Fish and Marine Sciences* 2012; 4(2): 159-64. doi: 10.5829/idosi.wjfm.s.2012.04.02.56406.
 26. Xu P, Zhang X, Wang X, Li J, Liu G, Kuang Y, et al. Genome sequence and genetic diversity of the common carp, *Cyprinus carpio*. *Nat Genet* 2014; 46(11): 1212-9. doi: 10.1038/ng.3098.
 27. Schultz S, Koussoroplis AM, Changizi-Magrhoor Z, Watzke J, Kainz MJ. Fish oil-based finishing diets strongly increase long-chain polyunsaturated fatty acid concentrations in farm-raised common carp (*Cyprinus carpio* L.). *Aquac Res* 2015; 46(9): 2174-84. doi: 10.1111/are.12373.
 28. Khan MN, Shahzad K, Chatta A, Sohail M, Piria M, Treer T. A review of introduction of common carp *Cyprinus carpio* in Pakistan: origin, purpose, impact and management. *Croatian Journal of Fisheries* 2016; 74(2): 71-80. doi: 10.1515/cjf-2016-0016.
 29. Macklin R, Brazier B, Harrison S, Chapman DV, Vilizzi L. A review of the status and range expansion of common carp (*Cyprinus carpio* L.) in Ireland. *Aquatic Invasions* 2016; 11(1): 75-82. doi: 10.3391/ai.2016.11.1.08.
 30. Sfakianakis DG, Renieri E, Kentouri M, Tsatsakis AM. Effect of heavy metals on fish larvae deformities: a review. *Environ Res* 2015; 137: 246-55. doi: 10.1016/j.envres.2014.12.014.
 31. Segner H, Schmitt-Jansen M, Sabater S. Assessing the impact of multiple stressors on aquatic biota: the receptor's side matters. *Environ Sci Technol* 2014; 48(14): 7690-6. doi: 10.1021/es405082t.
 32. Sinhohin VD, Sinhohin AP, Teixeira JM, Mileski KM, Hansen PC, Moreira PS, et al. Effects of the acute exposition to glyphosate-based herbicide on oxidative stress parameters and antioxidant responses in a hybrid Amazon fish surubim (*Pseudoplatystoma* sp.). *Ecotoxicol Environ Saf* 2014; 106: 181-7. doi: 10.1016/j.ecoenv.2014.04.040.
 33. Hoo JY, Kumari Y, Shaikh MF, Hue SM, Goh BH. Zebrafish: a versatile animal model for fertility research. 2016; 2016: 9732780. doi: 10.1155/2016/9732780.
 34. Montajami S, Hajiahmadyan M, Forouhar Vajargah M, Hosseini Zarandeh AS, Shirood Mirzaie F, Hosseini SA. Effect of symbiotic (*Bioimin imbo*) on Growth performance and survival rate of Texas cichlid (*Herichthys cyanoguttatus*) larvae. *Global Veterinaria* 2012; 9(3): 358-61. doi: 10.5829/idosi.gv.2012.9.3.6568.
 35. Forouhar Vajargah M, Hedayati A. Toxicity effects of cadmium in grass carp (*Ctenopharyngodon idella*) and Big Head Carp (*Hypophthalmichthys nobilis*). *Transylv Rev Syst Ecol Res* 2017; 19(1): 43-8. doi: 10.1515/trser-2017-0004.
 36. Bene C, Barange M, Subasinghe R, Pinstrup-Andersen P, Merino G, Hemre GI, et al. Feeding 9 billion by 2050 – Putting fish back on the menu. *Food Secur* 2015; 7(2): 261-74. doi: 10.1007/s12571-015-0427-z.
 37. Bajzelj B, Richards K, Allwood JM, Smith P, Dennis JS, Curmi E, et al. Importance of food-demand management for climate mitigation. *Nat Clim Chang* 2014; 4(10): 924-29. doi: 10.1038/nclimate2353.
 38. Sekhon BS. Nanotechnology in agri-food production: an overview. *Nanotechnol Sci Appl* 2014; 7: 31-53. doi: 10.2147/nsa.s39406.
 39. World Health Organization. World health statistics monitoring health for the SDGs, sustainable development goals. Geneva: WHO; 2016.
 40. Kruger M, Schledorn P, Schrod W, Hoppe HW, Lutz W, Shehata AA. Detection of glyphosate residues in animals and humans. *J Environ Anal Toxicol* 2014; 4(2): 1-5. doi: 10.4172/2161-0525.1000210.
 41. Kaur Gil IH, Garg H. Pesticides: environmental impacts

- and management strategies. In: Soloneski S, ed. Pesticides-Toxic Aspects. InTech; 2014. p. 188-230. doi: 10.5772/57399.
42. Akinsorotan AM. Toxicity of glyphosate herbicide on *Clarias gariepinus* fingerlings. Adv Res Biol Sci 2014; 2(1): 1-5.
 43. Micah AD, Adakole JA, Yusuf A, Mohammed NA. Acute effects of Glyphosate on the behavioural and hematological characteristics of hetero *Clarias* (hybrid) fingerlings. J Aquac Eng Fish Res 2017; 3(1): 13-18. doi: 10.3153/JAEFR17003.
 44. Henao Munoz LM, Montes Rojas CM, Bernal Bautista MH. Acute toxicity and sublethal effects of the mixture glyphosate (Roundup Active) and Cosmo-Flux 411F to anuran embryos and tadpoles of four Colombian species. Rev Biol Trop 2015; 63(1): 223-33.
 45. Szekacs I, Fejes A, Klatyik S, Takacs E, Patko D, Pomothy J, et al. Environmental and toxicological impacts of glyphosate with its formulating adjuvant. International Journal of Biological Veterinary Agricultural and Food Engineering 2014; 8(3): 219-24.
 46. Environmental Protection Agency. Ecological effects test guidelines; fish acute toxicity test, freshwater and marine. Washington, USA: United Environmental Protection Agency; 1996.