



Bioremediation of methyl tertiary-butyl ether (MTBE) by three pure bacterial cultures

Zabihollah Yousefi^{1*}, Zeinab Tahernezhad¹, Seyed Noroddin Mousavinasab², Reza Safari³, Ahmadrza Bekhradnia⁴

¹Department of Environmental Health Engineering, School of Public Health, Mazandaran University of Medical Sciences, Sari, Iran

²Department of Biostatistics, School of Public Health, Mazandaran University of Medical Sciences, Sari, Iran

³Iranian Fisheries Science Research Institute (IFSRI), Caspian Sea Ecology Research Center, Agricultural Research Education and Extension Organization (AREEO), Sari, Iran

⁴Department of Medicinal Chemistry, Pharmaceutical Sciences Research Center, Mazandaran University of Medical Sciences, Sari, Iran

Abstract

Background: Bioremediation of groundwater and soil contamination is more economical than physicochemical remediation. The present study focused on the bioremediation capability of two bacterial species (*Klebsiella planticola* and *Enterobacter cloacae*) from the family Enterobacteriaceae. These bacteria have been identified as new species with capability of degrading methyl tertiary-butyl ether (MTBE). In order to enhance their degradation capability, selected concentrations and retention time were investigated.

Methods: The bacteria were cultured on the nutrient agar (NA) medium at room temperature. pH of the medium was adjusted to 7. The medium was autoclaved at 121°C for 15 minutes and incubated for 24 hours at 35°C. After 24 hours, the mixture was inoculated into 50 mL of Luria Bertani (LB) liquid medium containing 50 and 150 ppm MTBE. The cultures were incubated for 2 and 5 days at 35°C and shaken on a shaker at 150 rpm. Cell concentrations of the bacteria in pure culture were determined from the optical density at 600 nm using a UV–VIS spectrophotometer. Then, the culture was centrifuged at 3800 rpm for 20 minutes. In the next step, the MTBE concentration in the supernatant was measured by gas chromatography/mass spectrometry (GC/MS, Agilent Technologies, 5975, US10304411, 5.02.07).

Results: The results showed that both strains are able to grow in the presence of 50 and 150 ppm MTBE. In the best conditions, when cell density was 3×10^8 CFU/mL during 5 days, the highest rate of MTBE degradation for *K. planticola* and *E. cloacae*, was 43% and 40%, respectively. It was also revealed that *Escherichia coli* can degrade 50 and 150 ppm MTBE about 19.8% and 13.65%, respectively.

Conclusion: It seems that *E. coli* can be a good candidate for MTBE degradation at high concentrations for a time longer than that in the present study. It was also found that the species have high performance at 50 ppm than 150 ppm. So, these bacteria can remove MTBE from the environment.

Keywords: Biodegradation, *Klebsiella planticola*, *Enterobacter cloacae*, *Escherichia coli*, methyl tertiary-butyl ether

Citation: Yousefi Z, Tahernezhad Z, Mousavinasab SN, Safari R, Bekhradnia A. Bioremediation of methyl tertiary-butyl ether (MTBE) by three pure bacterial cultures. Environmental Health Engineering and Management Journal 2018; 5(2): 123–128. doi: 10.15171/EHEM.2018.17.

Article History:

Received: 22 April 2018

Accepted: 27 May 2018

ePublished: 15 June 2018

*Correspondence to:

Zabihollah Yousefi

Email: zyousefi2004@gmail.com

Introduction

Unleaded gasoline consists hydrocarbons and different chemical compounds such as methyl tertiary-butyl ether (MTBE) (1,2). MTBE ($C_5H_{12}O$) is an oxygenate organic compound, that has been used as additive in gasoline since the late 1970s, to replace tetraethyl lead (TEL) and other toxic chemicals (3-5). MTBE is a persistent compound in the environment because it is highly soluble in water, poorly adsorbed by soil and is biologically and chemically stable against degradation (6). Accidental fuel leakage during storage or transportation is the main source of environmental contamination with MTBE (7).

Therefore, the presence of MTBE in water is responsible for taste and odor related issues, genotoxicity and skin and eye irritation. Taste and odor thresholds for MTBE are 20-40 ppb (8,9). Generally, MTBE is the most common oxygenate compound, because it is cost-effective and easy-to-use (7). Due to its economic benefits, bioremediation with 99% efficiency, is a more attractive option than physicochemical remediation technologies such as ozone utilization, activated carbon, vaporization extraction and other methods (3,7,10-12). All microorganisms are not able to degrade MTBE easily (13). Some bacteria such as *Pseudomonas*, *Rhodococcus*, *Mycobacterium*, *Enterobacter*,



and *Achromobacter* are capable of degrading MTBE co-metabolically but not tert-Butyl Alcohol (TBA) (1). A few bacterial strains such as *Methylibium petroleiphilum* PM1, *Hydrogenophaga flava* ENV735, *Achromobacter xylosoxidans* MCM 1/11, *Pseudomonas* sp., *Bacillus* sp., and *Streptococcus* sp. can utilize MTBE as the sole source of carbon and energy (14-18). Two bacterial isolates (IsoSL1 and Iso2A) degraded MTBE in both nutrient-rich and nutrient-limited media. The highest rate of MTBE degradation was reported 29.6% and 27.8%, respectively, in 28 days (19). Researchers reported that bacteria such as *A. xylosoxidans* MCM 1/11 can use MTBE in 7 days (17). In this study, bioremediation of MTBE by *Klebsiella planticola* and *Enterobacter cloacae* at laboratory conditions was investigated. On the other hand, according to the abundance of *Escherichia coli* and its capability to utilize a wide range of hydrocarbons while the engineered *E. coli* was used for bioremediation, therefore, MTBE-degrading capacity of this bacterium was also compared. MTBE is one of the gasoline components that, nowadays, is spreading in the environment and can pollute soil, water, and groundwater. The influence of microbial degradation of organic substances and MTBE is well known (17-19). Many studies have been conducted on the MTBE biodegradation by pure bacterial cultures such as *Bacillus cereus* and *Klebsiella terrigena*, *Enterobacter* sp. NKNUO2, and other microorganisms (7,14-17,20). The first step in bioremediation is selecting the best bacteria because only some bacteria can use MTBE as a source of carbon and energy. A few pure or mixed bacterial cultures can grow on MTBE and use it as a carbon and energy source and some strains grow slowly on this oxygenated compound with low cell yields (21). Due to the complex molecular structure of MTBE, this compound is resistant to biodegradation because its ether bond and tertiary carbon atom are relatively unreactive (21-23). The toxic effects of products produced during metabolism can cause it. The intermediate products of MTBE biodegradation are tert-butoxy methanol (TBM), formaldehyde and TBA, respectively (23-25). Steffan et al demonstrated that the growth rates of propane-oxidizing bacteria on MTBE is very slow. Numerous microorganisms including EVN 735, *Variovorax paradoxus* CL-8, *Chryseobacterium* sp. A-3, *B. cereus*, *K. terrigena*, *Enterobacter* sp., and NKNUO2, have also the capability to remove MTBE from the environment (12,20,26-30). The biological degradation of MTBE and most of the organic matter is nowadays known in science (31,32). In a study by Salanitro et al, biomass yields (gram of dry weight cells per gram of MTBE) were 0.21 to 0.28 (32). Some studies have also investigated MTBE biodegradation by *Mycobacterium* (33,34). But so far, no study has been conducted to investigate the role of *K. planticola* in bioremediation of MTBE, while some species of *Klebsiella* have been found to be capable of utilizing MTBE, n-hexadecane, and other hydrocarbons contaminating soil (20,35,36).

Materials and Methods

Materials

MTBE (GC purity $\geq 98\%$) was purchased from Persian Type Culture Collection (PTCC). Other chemicals were analytical grade and purchased from Merck (Darmstadt, Germany).

After sterilization and passing through a 2-mm mesh sieve, specific concentrations of MTBE were added to 10 g of soil, and the growth rate of microorganisms and the concentration of MTBE were measured.

Microorganisms and incubation conditions

In order to determine a strain capable of growing on MTBE, two concentrations and two retention time were examined. *K. planticola* and *E. cloacae* were purchased from the PTCC. Then, these bacteria were cultured on the nutrient agar (NA) medium at room temperature. The composition of NA was as follows (gr^{-1}): 0.5% Peptone, 0.3% beef extract/yeast extract, 1.5% agar, 0.5% sodium chloride, and distilled water. pH of the medium was adjusted to 7. The medium was autoclaved at 121°C for 15 minutes. In the next step, the cultures were incubated for 24 hours at 35°C. After 24 hours, the inoculum density at the beginning of the test were 1.5×10^8 CFU/mL and 3×10^8 CFU/mL, the mixture were then inoculated into 50 mL of Luria Bertani (LB) liquid medium containing 50 and 150 ppm MTBE (based on pretest) (12). LB medium was composed of (g/L): 10 g/L Trypton, 5 g/L yeast extraction, and 10 g/L NaCl. Then, the medium was autoclaved at 121°C for 15 minutes at pH 7. The cultures were incubated 2-5 days at 35°C in a shaker incubator at 150 rpm. Cell concentrations of *K. planticola* and *E. cloacae* in pure culture were determined from the optical density at 600 nm using a UV-VIS spectrophotometer (Hach model). Then, the culture was centrifuged in sealed tubes and cells were harvested from the medium by centrifugation at 3800 rpm for 20 minutes. Afterwards, the MTBE concentration in the supernatant was measured by gas chromatography/mass spectrometry (GC/MS, Agilent Technologies, 5975, US10304411, 5.02.07). The conditions for the GC analysis were as follows: 45°C held for 4 minutes, temperature ramped from 16°C/min to 70°C for 4.37 minutes, from 22°C/min to 100°C for 1.36 minutes, and 28°C/min to 220°C for 4.28 minutes (1,18,22). Helium was used as the carrier gas with an approximate flow rate of 1.10 mL/min. The sample without microorganisms was applied as blank in all tests. All specimens were tested two times. In this study, 38 samples were evaluated. Data were analyzed using statistical tests such as ANOVA, correlation, regression, and etc.

Results

During the incubation periods (2 and 5 days), *K. planticola* and *E. cloacae* were capable of growing on MTBE, as the source of carbon and energy, while initial concentrations of MTBE were studied. The removal rate of MTBE by two

Table 1. Biodegradation of MTBE by *Klebsiella planticola* and *Enterobacter cloacae*

Bacterial species	Initial concentration of MTBE (ppm)	Initial amount of bacteria (CFU/mL)	Time: 2 days		Time: 5 days	
			Removal percentage (%)	Residual value of concentration (ppm)	Removal percentage (%)	Residual value of concentration (ppm)
<i>K. planticola</i>	50	1.5×10 ⁸	11.8	44.1	19	40.5
		3×10 ⁸	29	35.5	43	21.5
	150	1.5×10 ⁸	6.95	139.5	12.96	130.5
		3×10 ⁸	24.3	12.5	33.5	99.5
<i>E. cloacae</i>	50	1.5×10 ⁸	9.1	45.45	16.3	41.85
		3×10 ⁸	24.6	37.7	40	30.05
	150	1.5×10 ⁸	5.3	142	11.9	132.15
		3×10 ⁸	15.95	126	30.95	103.5
<i>E. coli</i>	50	3×10 ⁸	24.3	12.5	33.5	99.5
		1.5×10 ⁸	9.1	45.45	16.3	41.85
	150	3×10 ⁸	24.6	37.7	40	30.05
		1.5×10 ⁸	5.3	142	11.9	132.15

pure bacterial cultures and the growth rate of two bacterial species at different concentrations of MTBE included in soil samples are shown in Table 1.

Biodegradation of 50 ppm MTBE by *K. planticola* and *E. cloacae* with inoculum sizes of 3×10⁸ CFU/mL, is displayed in Figure 1. As shown in Figure 1, *K. planticola* and *E. cloacae* respectively indicated 43% and 40% MTBE degradation, in 120 hours detention time. In addition, the bacteria showed 24.6% and 29% MTBE degradation after 48 hours.

Biodegradation of 150 ppm MTBE by *K. planticola* and *E. cloacae* with inoculum size of 3×10⁸ CFU/mL, is displayed in Figure 2. As shown in Figure 2, *K. planticola* and *E. cloacae* respectively indicated 30.95% and 33.50% MTBE degradation, in 120 hours detention time. The bacteria also showed 19.95% and 24.30% MTBE degradation after 48 hours.

48 hours.

Biodegradation of 150 ppm MTBE by *E. coli* with inoculum size of 3×10⁸ CFU/mL, is presented in Figure 3. As presented in Figure 3, *E. coli* with different inoculum sizes of 1.5×10⁸ and 3×10⁸ CFU/mL showed 13% and 19.8% MTBE degradation, respectively, in 120 hours detention time. The bacteria also showed 9% and 6.5% MTBE degradation after 48 hours.

Discussion

The bacteria were capable to grow in all samples while the growth rate of the bacteria and MTBE degradation rate were different. The results showed that both bacterial species, in the similar conditions, could use initial MTBE concentration of 50 ppm better than the concentration of 150 ppm. In a similar study by Okeke et al, concentration




Figure 1. Biodegradation of 50 ppm MTBE by *Klebsiella planticola* and *Enterobacter cloacae* with inoculum size of 3×10⁸ CFU/mL. White column, optical density of *K. planticola* after 48 and 120 hours and black column, optical density of *E. cloacae* after 48 and 120 hours.




Figure 2. Biodegradation of 150 ppm MTBE by *Klebsiella planticola* and *Enterobacter cloacae* with inoculum size of 3×10⁸ CFU/mL. White column, optical density of *K. planticola* after 48 and 120 hours and black column, optical density of *E. cloacae* after 48 and 120 hours.




Figure 3. Biodegradation of 150 ppm MTBE by *E. coli* with different inoculum sizes of 1.5×10^8 and 3×10^8 CFU/mL. White column, optical density of *E. coli* after 48 hours. .

of 50 ppm has been reported as the best concentration of MTBE. It seems that toxicity increases at higher concentrations (18). Also, in the study by Zhang et al, on the bioremediation of MTBE-contaminated soil, the results showed that MTBE degradation rate increased from 50 to below 100 ppm (12). Another study also showed that *Staphylococcus saprophyticus* 6sy and *Pseudomonas* sp. 24p were capable of degrading 25 $\mu\text{g/mL}$ MTBE better than 125 $\mu\text{g/mL}$ in 21 days. Other researchers insisted on the role of different concentrations of MTBE on bioremediation of MTBE (37). Abbaspour et al showed that *Bacillus cereus* strain RJ1 can survive at different concentrations of MTBE while the initial concentration of 200 ppm is rapidly degraded (38). In this study, the highest rate of MTBE degradation was recorded after 5 days while there were no significant MTBE disappearances in the blank samples. It was also revealed that pure culture of *K. planticola* is more capable of degrading MTBE in the initial concentration of 50 ppm comparing with *E. cloacae*, especially after the end of the period. *K. planticola* showed a significant difference (student's *t* test) in different concentrations ($P=0.005$) compared to *E. cloacae* ($P=0.0117$). The initial absorbance at 600 nm was 0.1 and the inoculum size was 3×10^8 CFU/mL. According to Figure 1, MTBE degradation rate by both strains increased with increasing the microbial density. The results are consistent with the results of the study by Zhang et al (12). They reported that there is a direct relationship between the removal of MTBE and time. According to Figures 1 to 3, all bacterial species had the potential for degradation of 50 ppm and 150 ppm MTBE. Biodegradation rate of MTBE was 29% and 43% while the absorbance at a wavelength of 600 nm was 0.315 and 0.367, respectively (Figure 1). *E. cloacae* is also a good candidate. Jose Barbera et al confirmed that

E. cloacae species play an important role in the MTBE biodegradation. They also showed that *E. cloacae* MCM2/1 has a high potential for utilizing MTBE (39). In the first hours, degradation rate by *K. planticola* was higher than other strains, and it seems that *E. coli* needs more time for degrading MTBE (Figure 3), it can be due to the complex molecular structure of MTBE, while the cultures containing yeast extract, this compound makes a good condition for better growth of the strains (19). Generally, there was no difference in the growth rate of both strains (*K. planticola* and *E. cloacae*) at different concentrations of MTBE. Besides, degradation rate was lower when the inoculum size was 1.5×10^8 CFU/mL (Table 1). The results of this study show that there is a direct relation between inoculum size and MTBE degradation. It should be noted that when the inoculum size increases, it uses more oxygen (12), which is consistent with the study by Rui-Ling et al. They reported that when cell density was 2×10^8 CFU/mL, the species had better performance than when it was 4.5×10^8 CFU/mL.

Conclusion

Klebsiella planticola, *E. cloacae* and *E. coli* could degrade MTBE at different concentrations and different cell densities in different times. Therefore, these strains are good candidates for removing MTBE from the environment. However, further studies using optimized media with other compounds for MTBE removal by these strains are suggested.

Acknowledgments

The authors would like to gratitude the Research Deputy of Mazandaran University of Medical Sciences for funding this study. And special thanks to P. Gil, Masoumeh Eslamifar and Masoumali Movahedi who helped us as research co-operators to perform this research.

Ethical issues

It is confirmed that this manuscript is the original work of the authors. The authors certify that all data collected during the study are presented in this manuscript, and no data from the study has been or will be published separately.

Competing interests

The authors declare that they have no conflicts of interests.

Authors' contribution

All authors contributed in data collection, analysis, and interpretation. All authors reviewed, refined, and approved the manuscript.

References

- Guisado IM, Purswani J, Gonzalez-Lopez J, Pozo C. Physiological and genetic screening methods for the isolation of methyl tert-butyl ether-degrading bacteria for

- bioremediation purposes. *Int Biodeterior Biodegradation* 2015; 97: 67-74. doi: 10.1016/j.ibiod.2014.11.008.
2. Schmidt TC, Haderlein SB, Pfister R, Forster R. Occurrence and fate modeling of MTBE and BTEX compounds in a Swiss Lake used as drinking water supply. *Water Res* 2004; 38(6): 1520-9. doi: 10.1016/j.watres.2003.12.027.
 3. Levchuk I, Bhatnagar A, Sillanpaa M. Overview of technologies for removal of methyl tert-butyl ether (MTBE) from water. *Sci Total Environ* 2014; 476-477: 415-33. doi: 10.1016/j.scitotenv.2014.01.037.
 4. The U.S. Environmental Protection Agency. MTBE in Fuels [cited 2017 Oct 23]. Available from: <http://www.epa.gov/mtbe/gas.htm>.
 5. Morse PM. Producers brace for MTBE Phaseout. *Chem Eng News* 1999; 77(15): 26-7. doi: 10.1021/cen-v077n015.p026.
 6. Soltani B, Moheb A. Ultrasound irradiation facilitated adsorption of MTBE from aqueous solution using exfoliated graphite. *Chem Eng Technol* 2010; 33(7): 1107-11. doi: 10.1002/ceat.200900345.
 7. Chen CS, Tien CJ, Zhan KV. Evaluation of intrinsic bioremediation of methyl tert-butyl ether (MTBE) contaminated groundwater. *Journal of Soil and Groundwater Environment* 2014; 19(5): 9-17. doi: 10.7857/JSGE.2014.19.5.009.
 8. Moreels D, Bastiaens L, Ollevier F, Merckx R, Diels L, Springael D. Evaluation of the intrinsic methyl tert-butyl ether (MTBE) biodegradation potential of hydrocarbon contaminated subsurface soils in batch microcosm systems. *FEMS Microbiol Ecol* 2004; 49(1): 121-8. doi: 10.1016/j.femsec.2004.02.016.
 9. Young WF, Horth H, Crane R, Ogden T, Arnott M. Taste and odour threshold concentrations of potential potable water contaminants. *Wat Res* 1996; 30(2): 331-40. doi: 10.1016/0043-1354(95)00173-5.
 10. Nau A, Kohl S, Zanthoff HW, Wiederhold H, Vogel H. Ozonized activated carbon as catalyst for MTBE-cleavage. *Appl Catal A Gen* 2011; 397: 103-111.
 11. U.S. Environmental Protection Agency. Technologies for Treating MTBE and Other Fuel Oxygenates. Washington, DC: Agency Office of Solid Waste and Emergency Response Office of Superfund Remediation and Technology Innovation; 2004.
 12. Zhang RL, Huang GQ, Lian JY, Li XG. Degradation of MTBE and TBA by a new isolate from MTBE-contaminated soil. *J Environ Sci (China)* 2007; 19(9): 1120-4.
 13. Davis LC, Erickson LE. A review of bioremediation and natural attenuation of MTBE. *Environmental Progress* 2004; 23(3): 243-52. doi: 10.1002/ep.10028.
 14. Nakatsu CH, Hristova K, Hanada S, Meng XY, Hanson JR, Scow KM, et al. *Methylibium petroleiphilum* gen. nov., sp. nov., a novel methyl tert-butyl ether-degrading methylotroph of the *Betaproteobacteria*. *Int J Syst Evol Microbiol* 2006; 56(Pt 5): 983-9. doi: 10.1099/ijls.0.63524-0.
 15. Streger SH, Vainberg S, Dong H, Hatzinger PB. Enhancing transport of hydrogenophaga flava ENV735 for bioaugmentation of aquifers contaminated with methyl tert-butyl ether. *Appl Environ Microbiol* 2002; 68(11): 5571-9.
 16. Lee EH, Cho KS. Effect of substrate interaction on the degradation of methyl tert-butyl ether, benzene, toluene, ethylbenzene, and xylene by *Rhodococcus* sp. *J Hazard Mater* 2009; 167(1-3): 669-74. doi: 10.1016/j.jhazmat.2009.01.035.
 17. Eixarch H, Constanti M. Biodegradation of MTBE by *Achromobacter xylosoxidans* MCM1/1 induces synthesis of proteins that may be related to cell survival. *Process Biochem* 2010; 45(5): 794-8. doi: 10.1016/j.procbio.2009.12.015.
 18. Okeke BC, Frankenberger WT Jr. Biodegradation of methyl tertiary butyl ether (MTBE) by a bacterial enrichment consortia and its monoculture isolates. *Microbiol Res* 2003; 158(2): 99-106. doi: 10.1078/0944-5013-00181.
 19. Makut MD, Ishaya P. Bacterial species associated with soils contaminated with used petroleum products in Keffi town, Nigeria. *Afr J Microbiol Res* 2010; 4(16): 1698-702.
 20. Nasrollahzadeh HS, Najafpour GD, Aghamohammadi N. Biodegradation of phenanthrene by mixed culture consortia in batch bioreactor using central composite face-centered design. *Int J Environ Res* 2007; 1(2): 80-87.
 21. Talaie AR, Jafaarzahe N, Talaie MR, Beheshti M. Biodegradation of aromatic compounds in crude oil by isolated microorganisms from environment. *J Zanjan Univ Med Sci* 2010; 18(70): 68-80. [In Persian].
 22. Arabi R, Bemanian SH, Taherzadeh MJ. Rapid biodegradation of Methyl tert - Butyl Ether (MTBE) by pure bacterial cultures. *Iran J Chem Chem Eng* 2007; 26(1): 1-7.
 23. Francois A, Mathis H, Godefroy D, Piveteau P, Fayolle F, Monot F. Biodegradation of methyl tert-butyl ether and other fuel oxygenates by a new strain, *Mycobacterium austroafricanum* IFP 2012. *Appl Environ Microbiol* 2002; 68(6): 2754-62.
 24. Smith CA, O'Reilly KT, Hyman MR. Characterization of the initial reactions during the cometabolic oxidation of methyl tert-butyl ether by propane-grown *Mycobacterium vaccae* JOB5. *Appl Environ Microbiol* 2003; 69(2): 796-804.
 25. HHatzinger PB, McClay K, Vainberg S, Tugusheva M, Condee CW, Steffan RJ. Biodegradation of Methyl tert -Butyl Ether by a pure bacterial culture. *Appl Environ Microbiol* 2001; 67(12): 5601-7. doi: 10.1128/AEM.67.12.5601-5607.2001
 26. Zaitsev GM, Uotila JS, Haggblom MM. Biodegradation of methyl tert-butyl ether by cold-adapted mixed and pure bacterial cultures. *Appl Microbiol Biotechnol* 2007; 74(5): 1092-102. doi: 10.1007/s00253-006-0737-3.
 27. Chen SC, Chen CS, Zhan KV, Yang KH, Chien CC, Shieh BS, et al. Biodegradation of methyl tert-butyl ether (MTBE) by *Enterobacter* sp. NKNU02. *J Hazard Mater* 2011; 186(2-3): 1744-50. doi: 10.1016/j.jhazmat.2010.12.079.
 28. Hanson JR, Ackerman CE, Scow KM. Biodegradation of methyl tert-butyl ether by a bacterial pure culture. *Appl Environ Microbiol* 1999; 65(11): 4788-92.
 29. Hatzinger PB, McClay K, Vainberg S, Tugusheva M, Condee CW, Steffan RJ. Biodegradation of methyl tert-butyl ether by a pure bacterial culture. *Appl Environ Microbiol* 2001; 67(12): 5601-7. doi: 10.1128/aem.67.12.5601-5607.2001.
 30. Steffan RJ, McClay K, Vainberg S, Condee CW, Zhang D. Biodegradation of the gasoline oxygenates methyl tert-butyl ether, ethyl tert-butyl ether, and tert-amyl methyl ether by propane-oxidizing bacteria. *Appl Environ Microbiol* 1997; 63(11): 4216-22.

31. Mo K, Lora CO, Wanken AE, Javanmardian M, Yang X, Kulpa CF. Biodegradation of methyl t-butyl ether by pure bacterial cultures. *Appl Microbiol Biotechnol* 1997; 47: 69–72.
32. Salanitro JP, Diaz LA, Williams MP, Wisniewski HL. Isolation of a bacterial culture that degrades methyl t-Butyl ether. *Appl Environ Microbiol* 1994; 60(7): 2593-6.
33. Fayolle F, Vandecasteele JP, Monot F. Microbial degradation and fate in the environment of methyl tert-butyl ether and related fuel oxygenates. *Appl Microbiol Biotechnol* 2001; 56(3-4): 339-49.
34. Alfonso-Gordillo G, Flores-Ortiz CM, Morales-Barrera L, Cristiani-Urbina E. Biodegradation of methyl tertiary butyl ether (MTBE) by a microbial consortium in a continuous up-flow packed-bed biofilm reactor: kinetic study, metabolite identification and toxicity bioassays. *PLoS One* 2016; 11(12): e0167494. doi: 10.1371/journal.pone.0167494.
35. Nduka JK, Umeh LN, Okerulu IO, Umedum LN, Okoye HN. Utilization of different microbes in bioremediation of hydrocarbon contaminated soils stimulated with inorganic and organic fertilizers. *J Pet Environ Biotechnol* 2012; 3(2): 1-9. doi: 10.4172/2157-7463.1000116.
36. Dutra ES, Pascon RC, Vallim MA. Sao Paulo Zoo composting as a source of bacteria with bioremediation potential. *Afr J Microbiol Res* 2013; 7(45): 5200-6. doi: 10.5897/AJMR2013.5874.
37. Lalevic B, Raicevic V, Kikovic D, Jovanovic L, Surlan-Momirovic G, Jovic J, et al. Biodegradation of MTBE by bacteria isolated from oil hydrocarbons- contaminated environments. *Int J Environ Res* 2012; 6(1): 81-6. doi: 10.22059/ijer.2011.474.
38. Abbaspour M, Javid AH, Jalilzadeh Yengjeh R, Hassani AH, Ghavam Mostafavi P. The Biodegradation of methyl tert-Butyl ether (MTBE) by indigenous *Bacillus cereus* strain RJ1 isolated from soil. *Petroleum Science and Technology* 2013; 31(18): 1835-41. doi: 10.1080/10916466.2011.611562.
39. Jose Barbera M, Mateo E, Monkaityte R, Constanti M. Biodegradation of methyl tert-butyl ether by newly identified soil microorganisms in a simple mineral solution. *World J Microbiol Biotechnol* 2011; 27(4): 813-21. doi: 10.1007/s11274-010-0522-4.