








The concentration and human health risks attributed to pesticide residues in cucumber: A systematic review and meta-analysis

Pouria Karimi¹, Sepideh Sadeghi^{1,2}, Fatemeh Kariminejad³, Negin Rezaeiashad², Yadolah Fakhri⁴, Mohsen Sadani^{2*}, Farshid Soleimani⁵, Mohammad Sadeghi⁶

¹Student Research Committee, School of Public Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²Department of Environmental Health Engineering, School of Public Health and Safety, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³Student Research Committee, Department of Environmental Health Engineering, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

⁴Food Health Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

⁵Tobacco and Health Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

⁶Student Research Committee, School of Public Health, North Khorasan University of Medical Sciences, Bojnurd, Iran

Abstract

Background: Prolonged consumption of vegetables containing pesticide residues can pose a risk to the health of individuals over time.

Methods: This study aimed to retrieve the studies on the concentration of pesticides such as acetamiprid, chlorpyrifos, and diazinon in cucumber through a systematic review, and then, a meta-analysis focusing on specific subgroups. Finally, the hazard quotient (HQ) was used to assess the non-carcinogenic risk posed to consumers.

Results: The sequence of pesticides according to pooled (mean) concentration was acetamiprid (140.91 µg/kg) > diazinon (59.03 µg/kg) > chlorpyrifos (40.49 µg/kg). Also, the countries were sorted based on the pooled concentration of chlorpyrifos: Spain (180.00 µg/kg) > Egypt (124.90 µg/kg) > Kazakhstan (49.95 µg/kg) > Greece (20.50 µg/kg) > Saudi Arabia (20.00 µg/kg) > Jordan (3.54 µg/kg) > Iran (3.10 µg/kg). The order for diazinon and acetamiprid was Iran (43.20 µg/kg) > Saudi Arabia (33.00 µg/kg) > China (4.60 µg/kg) and Turkey (231.70 µg/kg) > Egypt (220 µg/kg) > Pakistan (100.00 µg/kg) > Kazakhstan (99.34 µg/kg), respectively.

Conclusion: The HQ of acetamiprid, chlorpyrifos, and diazinon was less than 1 for all countries; hence, the consumption of cucumbers containing these pesticides does not pose a non-carcinogenic risk.

Keywords: Agrochemical, Exposure, Meta-analysis, Pesticide, Risk assessment

Citation: Karimi P, Sadeghi S, Kariminejad F, Rezaeiashad N, Fakhri Y, Sadani M, et al. The concentration and human health risks attributed to pesticide residues in cucumber: a systematic review and meta-analysis. Environmental Health Engineering and Management Journal. 2025;12:1372. doi: 10.34172/EHEM.1372.

Article History:

Received: 15 May 2024

Accepted: 26 October 2024

ePublished: 7 January 2025

*Correspondence to:

Mohsen Sadani,
Email: m.sadani82@outlook.com

Introduction

In response to the growing global population, food producers must increase food production and minimize waste to ensure an adequate food supply (1,2). This is essential to meet the rising demand for food by scaling up production and increasing crop yield per unit of area (3,4). According to the United Nations, it is estimated that the world population will reach 9.8 billion by 2050 (5). The Food and Agriculture Organization (FAO) states that meeting the nutritional needs of the global population in 2050 would require increasing total food production by approximately 70% (6,7).

Agrochemicals, including fertilizers and pesticides, have

become essential in modern agriculture for increasing yields with less effort. Additionally, these nutrients promote crop growth and soil health (8). About 33% of global crop productivity could be compromised by pests such as diseases, rodents, weeds, plant-feeding insects, and mites. Pesticides are the most common substances used to control these detrimental elements (9,10). They are biological and chemical substances extensively utilized in modern agriculture to enhance crop yields by safeguarding plants through methods such as deterring, incapacitating, eliminating, or discouraging pests (11,12).

Organophosphate pesticides (OPs) can have harmful impacts on human health. Approximately 0.1% of the



pesticides applied reach the intended pests, while the rest are dispersed into water, soil, and food (13). The prevalence of pesticide poisoning among humans worldwide has risen significantly. In 1972, there were about 500 000 cases per year, which increased to approximately 25 million cases per year in 1990 (14). Pesticides pose significant risks to human health, affecting various bodily systems (12). They can disrupt metabolic homeostasis and with other variables have been associated with disturbances in the endocrine system, insulin resistance, nonalcoholic fatty liver disease, obesity, diabetes, blood cancer, and increased risks of lung and prostate cancer (15). Exposure to OPs encompasses more than just occupational contact, extending to contact through dust and aerosols, as well as the ingestion of contaminated food (16,17).

Food security is a significant public concern worldwide (18,19). The FAO defines food security as the physical, social, and economic access of all individuals to adequate, safe, and nutritious food at all times, ensuring a healthy and active life (20). The United States Environmental Protection Agency (USEPA) classifies the majority of OPs based on their toxicity levels for inhalation and oral exposure, using classes I to IV (21).

Vegetables and fruits are essential elements of diets promoted under the term “healthy lifestyle” because they offer vital nutrients, dietary fiber, and vitamins. According to the World Health Organization (WHO), vegetables and fruits typically account for approximately 30% of food consumption worldwide (22,23). The existence of organophosphate residues in vegetables and fruits could endanger consumer health, as these substances are a significant component of the human diet (24,25).

Cucumber (*Cucumis sativus* L.) is a commonly cultivated crop from the Cucurbitaceae family, grown in open fields and protected houses (26). It is a healthy fruit with numerous benefits, including maintaining blood pressure, regulating hydration, controlling blood sugar, aiding in weight loss, and promoting skin health and digestion. Cucumbers are high in water content, vitamins (K, C, A), and minerals (potassium, manganese, magnesium), and they also contain fiber. Additionally, they have hydrating, antimicrobial, and detoxifying properties, and may help prevent cancer and bone diseases (27,28).

To this end, several studies have been conducted globally to evaluate the presence of pesticide residues in various vegetables. For instance, in 2018, a total of 19 samples of cucumber and tomato were randomly collected from 14 greenhouse farms. The findings showed that diazinon, malathion, chlorpyrifos, ethephon, profenofos, and oxyfluorfen were detected in 94.7% of the samples. The residue levels in the cucumber samples varied between 0.05 and 167.19 mg/kg, while those in the tomato samples ranged from 2.55 to 136.87 mg/kg, all exceeding the maximum residue limits (MRLs) established by the Codex Alimentarius (29). In another

study conducted in Iran, 64 samples of fresh cucumbers were analyzed for the presence of four pesticides (diazinon, imidacloprid, Pirimicarb, and acetamiprid). The cucumbers were randomly sampled 2, 5, 10, and 14 days after pesticide application. The results showed that acetamiprid and Pirimicarb had higher persistence in cucumbers compared to the other two pesticides. The lowest residue levels in cucumbers were detected 14 days after the application of acetamiprid (2.06 mg/kg) and pirimicarb (2.12 mg/kg). In contrast, the lowest residue levels of diazinon and imidacloprid were 0.24 and 1.16 mg/kg, respectively, within 14 days. All tested residues dissipated completely 21 days after application in cucumbers (30). Likewise, another investigation in Malaysia examined the possible presence of diazinon, chlorpyrifos, and dimethoate residues in cucumbers and tomatoes conveniently sampled from four food outlets within Taman University. The research found that there were no traces of diazinon, chlorpyrifos, and dimethoate residues in the cucumber and tomato samples collected from these permanent food outlets (31).

Despite existing concerns and evidence about the unprincipled use of pesticides in agriculture, the effects related to human health have not been comprehensively investigated so far. In the present study, cucumber was considered as a product that is consumed all over the world. This study is a systematic review and meta-analysis that addresses a significant gap in previous studies by conducting a thorough review and analysis of pesticide levels in cucumbers, aiming to understand their potential impact on human health. To achieve this, the study specifically focuses on examining the concentrations of chlorpyrifos, acetamiprid, and diazinon in cucumber, a widely used vegetable in diverse cuisines worldwide. By synthesizing data from various studies, this review takes a meticulous approach to evaluate the overall extent of pesticide contamination in cucumbers and understand the non-carcinogenic health risks associated with consuming them. The results of this investigation will greatly enhance our comprehension of the current state of pesticide residues in cucumbers. By illuminating this issue, the study will provide essential information that can be employed to develop efficient measures to reduce these risks and safeguard the food supply for consumers.

Materials and Methods

Search strategy

The literature search was carried out following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Figure 1) (32,33) through scientific databases such as PubMed, Web of Science, and Scopus from January 1, 2010 to April 1, 2022. The search was based on the following terms: “Cucumber OR *Cucumis sativus*” OR *C. sativus*” AND “pesticide OR acetamiprid OR chlorpyrifos OR diazinon”.




Figure 1. The selection process of articles based on the PRISMA protocol

Disagreements in the selection of articles were resolved through the final comment of the correspondence (34).

Inclusion and exclusion criteria

Among the mentioned databases, the studies that evaluated pesticide levels in vegetables were included based on the following criteria: The articles were published in English, the articles that their full text was accessible, the original studies distinctly monitored the concentration of chlorpyrifos and diazinon in cucumber, and those that reported statistical details (mean, standard deviation [SD], and/or standard error [SE]) about the pesticides. In addition, the exclusion criteria were: Theses, clinical studies, abstracts, books, case series, case reports, conference papers, review studies, and letters to the editor; the lab-studies that estimated the fate and behavior of pesticides in water-soil-plant, and finally, lab-studies that evaluated the analytical methods.

Data extraction

In this research, the collected data included the author(s) name, publication year, place (country) of sampling, type of pesticide, statistical details (mean, SD/ or SE, and sample size), and title of the study. After the extraction of data, the statistical details about the concentration of

the chemicals were unified as mg/kg (wet-weight) of the vegetable.

Statistical and meta-analysis of data

The meta-analysis of the extracted data was conducted using STATA version 14.0 (Stata Corp, College Station, TX). After unifying the data units, mean concentration, and SE were considered for the analysis. SE was calculated using SD and sample size (N) according to Eq. 1(35).

$$SE = \frac{SD}{\sqrt{N}} \quad (1)$$

The subgroup analysis was conducted separately according to the sampling country. Heterogeneity, as an index describing the properties of a dataset, was assessed using I^2 . According to Cochrane recommendations, I^2 higher than 50% indicates substantial heterogeneity. Since the dataset obtained in this study showed heterogeneity greater than 90%, an analysis was conducted using a random effects model (36).

Health risk assessment

The estimation of exposure among the investigated countries, as the initial stage in evaluating health risk was

conducted by Eq. (2) (37).

$$EDI = \frac{C_m \times IR \times EF \times ED}{BW \times ATn} \quad (2)$$

Here, *EDI* represents the estimated daily intake (mg/kg.day), C_m is the concentration of pesticides (mg/kg-wet weight), *IR* is the ingestion rate (g/n-d), *EF* is the exposure frequency (350 day/year), *ED* is the exposure duration (children=6 years and adults=30 years), *BW* is body weight (children=15 kg and adults=70 kg), and *ATn* ($365 \times ED$) is average time exposure (children=2190 days and adults=10950 days). The ingestion rate for cucumber was assumed to be 10% of the human vegetable ingestion rate obtained from the Food and Agriculture Organization of the United Nations (FAO) statistics (38).

To evaluate the hazard quotient (HQ) in association with non-carcinogenic risk (n-CR), Eq. (3) was used. Where, *RfD* implies reference dose, which is considered to be 0.001, 0.0007, and 0.07 for chlorpyrifos, diazinon, and acetamiprid, respectively (39,40). The cumulative risk associated with the simultaneous ingestion of all chemicals is calculated via Eq. (4) (41).

$$HQ = \frac{EDI}{RfD} \quad (3)$$

$$THQ = \sum_{i=1}^n HQ_i \quad (4)$$

Results

Selection process and characteristics of studies

The present study primarily focused on the systematic review and meta-analysis of the concentration of acetamiprid, chlorpyrifos, and diazinon in cucumbers. This involved searching through the databases Scopus, Web of Science, and PubMed, resulting in the retrieval of 2767 articles for analysis. After the removal of 1424 duplicate papers, the titles and abstracts of the selected

studies were screened for eligibility, leading to the exclusion of 1280 articles that did not meet the criteria. Ultimately, 20 articles were included that fulfilled all required criteria (Figure 1).

Of the articles finally selected, which were conducted in 11 countries (Turkey, Egypt, Kazakhstan, Pakistan, Greece, Jordan, Iran, Saudi Arabia, Kazakhstan, Spain, and China), the ranking of years based on the number of studies conducted in was as follows: 2014 > 2016 > 2020 ≈ 2021 (Figure 2-a). Most of these studies focused on monitoring chlorpyrifos (41%) ≈ diazinon (41%) compared to acetamiprid (18%) (Figure 2-b), which can be due to the higher consumption of Chlorpyrifos and Diazinon in agriculture and household pest control (42,43).

It should be noted that the usage amounts of these pesticides may vary depending on factors such as crop types, pest infestations, and regional preferences, and can differ from one country to another (44). The characteristics of the included articles are presented in Table 1.

Meta-analysis concentration of pesticides

The concentration of pesticides

According to the results indicated in Tables 2-4, the overall ranking of countries based on the concentration of chlorpyrifos was as follows: Spain (180.00 µg/kg) > Egypt (124.90 µg/kg) > Kazakhstan (49.95 µg/kg) > Greece (20.50 µg/kg) > Saudi Arabia (20.00 µg/kg) > Jordan (3.54 µg/kg) > Iran (3.10 µg/kg). The order for diazinon and acetamiprid was Iran (43.20 µg/kg) > Saudi Arabia (33.00 µg/kg) > China (4.60 µg/kg) and Turkey (231.70 µg/kg) > Egypt (220 µg/kg) > Pakistan (100.00 µg/kg) > Kazakhstan (99.34 µg/kg), respectively. The obtained results from the present study, reported in Tables 2-4, showed the relative ranking of pesticides based on their concentrations in cucumber as follows: Acetamiprid (140.91 µg/kg) > Diazinon (59.03 µg/kg) > Chlorpyrifos (40.49 µg/kg).

Health risk assessment

According to the results reported in Table 5, the HQ




Figure 2. Statistical information about the included articles: (a) the number of selected articles reported per year, (b) the contribution of studies by the type of pesticide

Table 1. The characteristics of the included articles

Reference	Author	Year	Country	Unit	Pesticides	Sample Size	Mean	SD	SE
(45)	Yu et al	2016	China	µg/kg	Diazinon	31	4.600	0.506	0.091
(46)	Vlassi et al	2020	Greece	µg/kg	Chlorpyrifos	10	20.500	2.255	0.713
(47)	Ibrahim et al	2019	Jordan	µg/kg	Chlorpyrifos	5	7.000	30.000	13.416
(47)	Ibrahim et al	2019	Jordan	µg/kg	Chlorpyrifos	5	3.536	0.389	0.174
(47)	Ibrahim et al	2019	Jordan	µg/kg	Chlorpyrifos	5	3.536	0.389	0.174
(47)	Ibrahim et al	2019	Jordan	µg/kg	Chlorpyrifos	5	3.536	0.389	0.174
(47)	Ibrahim et al	2019	Jordan	µg/kg	Chlorpyrifos	5	3.536	0.389	0.174
(48)	Sungur & Tunur	2012	Turkey	µg/kg	Acetamiprid	10	437.000	20.000	6.325
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	12	13.300	0.720	0.208
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	12	10.420	0.760	0.219
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	10	9.260	1.070	0.338
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	6	17.200	3.500	1.429
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	12	13.770	2.100	0.606
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	5	9.550	0.540	0.241
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	6	10.040	0.980	0.400
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	11	9.840	1.210	0.365
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	5	9.210	1.130	0.505
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	10	18.400	3.340	1.056
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	16	11.540	2.070	0.518
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	7	10.870	1.070	0.404
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	5	12.090	1.310	0.586
(49)	Shokrzadeh et al	2017	Iran	µg/kg	Diazinon	5	9.200	1.103	0.493
(50)	Shalaby et al	2021	Egypt	µg/kg	Acetamiprid	16	220.000	24.200	6.050
(50)	Shalaby et al	2021	Egypt	µg/kg	Chlorpyrifos	16	220.000	24.200	6.050
(51)	Nasiri et al	2016	Iran	µg/kg	Chlorpyrifos	43	0.097	0.001	0.000
(52)	Mosleh et al	2014	Saudi Arabia	µg/kg	Chlorpyrifos	5	20.000	2.000	0.894
(53)	Lozowicka et al	2015	Kazakhstan	µg/kg	Acetamiprid	19	250.000	27.500	6.309
(53)	Lozowicka et al	2015	Kazakhstan	µg/kg	Acetamiprid	19	10.000	1.100	0.252
(53)	Lozowicka et al	2015	Kazakhstan	µg/kg	Acetamiprid	19	100.000	11.000	2.524
(53)	Lozowicka et al	2015	Kazakhstan	µg/kg	Acetamiprid	19	10.000	1.100	0.252
(53)	Lozowicka et al	2015	Kazakhstan	µg/kg	Acetamiprid	19	150.000	16.500	3.785
(53)	Lozowicka et al	2015	Kazakhstan	µg/kg	Chlorpyrifos	19	70.000	7.700	1.767
(53)	Lozowicka et al	2015	Kazakhstan	µg/kg	Chlorpyrifos	19	30.000	3.300	0.757
(53)	Lozowicka et al	2015	Kazakhstan	µg/kg	Chlorpyrifos	19	50.000	5.500	1.262
(54)	Jamshidi et al	2016	Iran	µg/kg	Diazinon	106	4850.000	7440.000	722.637
(54)	Jamshidi et al	2016	Iran	µg/kg	Diazinon	21	10.000	30.000	6.547
(54)	Jamshidi et al	2016	Iran	µg/kg	Diazinon	85	6560.000	8320.000	902.431
(55)	Ganjeizadeh Rohani et al	2014	Iran	µg/kg	Diazinon	17	264.000	29.040	7.043
(55)	Ganjeizadeh Rohani et al	2014	Iran	µg/kg	Diazinon	17	832.000	91.520	22.197
(55)	Ganjeizadeh Rohani et al	2014	Iran	µg/kg	Diazinon	17	1113.000	122.430	29.694
(55)	Ganjeizadeh Rohani et al	2014	Iran	µg/kg	Diazinon	17	173.000	19.030	4.615
(55)	Ganjeizadeh Rohani et al	2014	Iran	µg/kg	Diazinon	17	526.000	57.860	14.033
(55)	Ganjeizadeh Rohani et al	2014	Iran	µg/kg	Diazinon	17	582.000	64.020	15.527
(56)	EL-Saeid & Selim	2013	Saudi Arabia	µg/kg	Diazinon	1057	33.000	3.630	0.112
(57)	Davoodi et al	2014	Iran	µg/kg	Diazinon	10	264.000	295.000	93.287
(58)	Darvishnejad et al	2020	Iran	µg/kg	Chlorpyrifos	17	3.100	0.341	0.083
(58)	Darvishnejad et al	2020	Iran	µg/kg	Diazinon	17	3.400	0.374	0.091

Table 1. Continued.

Reference	Author	Year	Country	Unit	Pesticides	Sample Size	Mean	SD	SE
(59)	Bazmandegan-Shamili et al	2017	Iran	µg/kg	Diazinon	17	1.510	0.060	0.015
(60)	Bakırcı et al	2014	Turkey	µg/kg	Acetamiprid	69	26.500	2.915	0.351
(61)	Bagheri et al	2012	Iran	µg/kg	Diazinon	7	55.000	5.100	1.928
(61)	Bagheri et al	2012	Iran	µg/kg	Diazinon	7	82.000	6.900	2.608
(62)	Ahmed et al	2014	Egypt	µg/kg	Chlorpyrifos	17	30.000	2.000	0.485
(63)	Quijano et al	2016	Spain	µg/kg	Chlorpyrifos	25	180.000	19.800	3.960
(64)	Siddique et al	2021	Pakistan	µg/kg	Acetamiprid	17	100.000	29.000	7.034

Table 2. Meta-analysis concentration of chlorpyrifos in cucumber (µg/kg)

Study	Number of studies	ES ^a	Lower	Upper	Weight (%)	Heterogeneity Statistic	Degrees of freedom	P value	I ²
Greece	1	20.500	19.103	21.897	7.93	0.00	0	.	0.0%
Jordan	1	3.536	3.366	3.707	33.88	0.07	4	0.999	0.0%
Egypt	2	124.904	0	311.101	12.78	979.97	1	0.000	99.9%
Iran	2	3.100	2.937	3.263	8.00	0.00	0	.	0.0%
Saudi Arabia	1	20.000	18.248	21.752	7.89	0.00	0	.	0.0%
Kazakhstan	1	49.955	27.867	72.044	23.28	524.64	2	0.000	99.6%
Spain	1	180.000	172.239	187.761	6.23	0.00	0	.	0.0%
Overall	9	40.489	36.368	44.610	100.00	11004.82	13	0.000	99.9%

^a Effect size: Weighted average.

Table 3. Meta-analysis concentration of diazinon in cucumber (µg/kg)

Study	Number of studies	ES ^a	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	P value	I ²
China	1	4.600	4.422	4.778	4.31	0.00	0	.	0.0%
Iran	7	43.204	40.238	46.170	91.38	20246.48	27	0.000	99.9%
Saudi Arabia	1	33.000	32.780	33.220	4.31	0.00	0	.	0.0%
Overall	9	59.030	54.586	63.473	100.00	97222.10	29	0.000	100.0%

^a Effect size: Weighted average.

Table 4. Meta-analysis concentration of acetamiprid in cucumber (µg/kg)

Study	Number of studies	ES ^a	Lower	Upper	Weight (%)	Heterogeneity statistic	Degrees of freedom	P value	I ²
Turkey	2	231.701	0	633.985	22.26	4199.23	1	0	100.00%
Egypt	1	220	208.142	231.858	10.72	0	0	.	0.0%
Kazakhstan	1	99.342	85.132	113.552	56.58	4058.48	4	0	99.90%
Pakistan	1	100	86.214	113.786	10.44	0	0	.	0.0%
Overall	5	140.912	126.899	154.926	100	11392.4	8	0	99.90%

^a Effect size: Weighted average.

values of all three types of pesticides for both children and adults were less than 1 in all countries, implying no appreciable non-cancer health risk.

Discussion

The consumption of pesticides varies across countries due to a range of factors, including agricultural methods, the social processes of pesticide application, economic factors, food security regulations, and politics (65,66). Although some developed countries have made noticeable progress

in decreasing pesticide use intensity, there has been a significant increase in pesticide usage in some developing countries. This has resulted in an uneven geographical distribution of pesticide consumption (67). However, the reasons for the high consumption of pesticides in countries with the highest consumption levels have been investigated.

The California red scale (*Aonidiella aurantii*) is the main citrus pest in Spain (68). Chlorpyrifos, an insecticide, has been the foundation for managing red-scale in Spain for

Table 5. Non-carcinogenic HQ risk in adults and children due to the ingestion of cucumbers containing pesticides

Age group	Country	Chlorpyrifos	Diazinon	Acetamiprid
Adult	China	-	0.008	-
	Greece	0.012	-	-
	Jordan	0.001	-	-
	Egypt	0.038	-	0.001
	Iran	0.001	0.019	-
	Saudi Arabia	0.003	0.007	-
	Kazakhstan	0.036	-	0.001
	Pakistan			0.000
	Spain	0.034	-	-
	Turkey			0.001
Child	China	-	0.037	-
	Greece	0.054	-	-
	Jordan	0.004	-	-
	Egypt	0.176	-	0.004
	Iran	0.005	0.089	-
	Saudi Arabia	0.014	0.032	-
	Kazakhstan	0.166	-	0.004
	Pakistan			0.000
	Spain	0.160	-	-
	Turkey			0.006

many years (69). This highly effective active ingredient is fast-acting, providing a quick 'knock-down' effect on pests (70). Due to the lack of an alternative to chlorpyrifos with this activity profile, its availability and consumption are very high in Spain, leading to increased usage across all types of crops (71). For example, chlorpyrifos was detected in 22% of samples such as apples, carrots, bananas, citrus fruits, grapes, lettuce, peaches, pears, potatoes, tomatoes, and cucumbers in Valencia (Spain) (63).

There are different conditions in Egypt due to the country's heavy reliance on agriculture and the use of pesticides to protect crops from pests and diseases (72). In this area, improper use of agricultural pesticides has increased the pollution of the Nile River, the main source of irrigation (73). As a result, irrigation with this contaminated water can lead to contamination of soil and agricultural products, as reported by many Egyptian studies (74-76). Among the different pesticides used in Egypt, Chlorpyrifos is one of the most frequently utilized OPs (77,78). It is consumed at approximately 1280 tons each year, according to statistics from the Agricultural Pesticides Committee, Ministry of Agriculture & Land Reclamation (79). Similar studies in local markets of Egypt have shown that Chlorpyrifos is commonly found alongside at least one other compound in soil and crop samples from this area, and residues of this pesticide were detected in three types of crops (74). Although its application is government-regulated, the precise dates of

application differ among fields depending on local insect infestations (80). Ibrahim et al, mentioned that increasing awareness among Egyptian farmers to reduce pesticide usage should be achieved through training on reasonable usage, following instructions on information labels, and complying with estimated safety periods (81).

Diazinon, an organophosphorus pesticide, is extensively used to control aphids, caterpillars, moths, butterflies, various worms, locusts, grasshoppers, and scales in pastures, orchards, vegetables, and field crops due to its high effectiveness and medium-low toxicity (82,83).

It is widely used in Iranian agriculture due to its low cost and effectiveness (84,85). According to a study (86) in Iran during 2012–2014, approximately 14000 tons of pesticides were used, with Diazinon making up 5% of the total amount. Although the majority of Diazinon's use in Iran is related to the northern regions and agricultural lands (87), some studies indicate its application in other areas as well. For example, Ebrahimzadeh et al study., conducted in Sistan and Baluchistan province (south-east of Iran), Diazinon is extensively utilized for agricultural purposes (88). The researchers mentioned that due to the increased use of Diazinon in spring, the season marking the beginning of crop cultivation, and the higher rainfall in this season, there is a higher probability of Diazinon transfer to the soil, and ultimately, more absorption in crops (89). Furthermore, the lack of a modern irrigation system in Iran has been identified as another factor contributing to the pesticide's transfer to water sources. Creating a network that integrates traditional knowledge of pests with scientific pesticide knowledge, sharing innovations that mitigate the adverse effects of pesticides, providing farmers with information that encourages changes in harmful pesticide practices, establishing close communication between agricultural experts and farmers, and updating the technical knowledge of agricultural experts in the management and usage of pesticides are the most important policy measures to improve the effective use of pesticides (90).

In Saudi Arabia, along with the expansion of agricultural land, the consumption of pesticides has also increased (91). This country was among the early adopters of the Stockholm Convention in 2002. However, due to the lack of data on persistent organic pollutants, it has not fully ratified the agreement (92). Meanwhile, Diazinon was the most commonly found pesticide in this area (93). Case in point, in a study conducted in the Asir Region of Saudi Arabia, 41.7% of Cucumber samples had detectable pesticide residues above the MRLs (94). The increasing pest attacks on important crops have been a significant issue in recent years (95). Additionally, climate change has exacerbated the incidence and spread of plant pests and diseases in this region, making crops more susceptible to various pests and diseases, thereby impacting crop productivity in recent years (96). This

rise in pest populations necessitates the implementation of control and monitoring programs (97). Reducing government involvement in distribution to decrease the need for maintaining large pesticide stocks, encouraging private sector participation, controlling prices, reducing subsidies or preferential tariffs on pesticides, limiting pesticide stock, and establishing strict regulation and control systems can help prevent the residues of pesticides in Saudi Arabia.

Acetamiprid is a neonicotinoid insecticide widely used on a variety of crops to control insect pests such as aphids, whiteflies, and beetles (98). It acts as a nicotinic acetylcholine receptor (nAChR) agonist, making it highly efficient in pest control (99). Turkey applies 33 million kg of pesticides annually, with 29% consisting of insecticides like Acetamiprid (100). The most important reason for the widespread use of Acetamiprid in Turkish agriculture is its effectiveness in controlling insects such as aphids, which can damage leafy plants, resulting in residues that can pose a threat to human health (101). The use of Acetamiprid in Turkey is regulated by the Turkish Food Codex and the European Union's maximum acceptable limits (102). However, there have been instances where residues of Acetamiprid were detected in cucumber samples at concentrations exceeding the maximum acceptable limit (103). In a study (104) conducted in supermarkets, bazaars, and greengrocer shops across Turkey, about 30.77% of the samples contained Acetamiprid residue above the MRL value (105,106). It was mentioned that some of the reasons for the high use of this pesticide in Turkey are its effectiveness in controlling pests, availability, and regulatory compliance, making it a convenient choice for farmers and agricultural workers. Meanwhile, over 50% of Turkish farmers and agricultural workers believe that pesticides are not harmful to them. Therefore, it is essential for governmental agencies, as well as the private sector, to address these issues through educational strategies (107).

According to the literature, the most important actions to decrease pesticide usage, improve their management, and subsequently, reduce their harmful effects in various countries include increasing awareness and training, researching and developing alternative pest management techniques and non-chemical methods, and implementing policies and regulations that promote sustainable and reduced pesticide use. It should be noted that the effectiveness of methods to reduce pesticide use varies across countries due to a complex interplay of agricultural, regulatory, economic, cultural, and technological factors.

The concentration of these pesticides in cucumbers can be influenced by various factors, such as surface evaporation, which depends on temperature conditions, as well as chemical or biochemical decomposition, metabolism, and photolysis (108). These results clearly show that

Acetamiprid is used more frequently in cucumbers compared with the other two pesticides. One reason may be related to the high availability, and subsequently, improper use or overuse of it (109). As mentioned earlier, Acetamiprid, a neonicotinoid insecticide, poses lower risks to mammals and the environment in comparison with organophosphate insecticides like diazinon and chlorpyrifos (98,110). Previous studies have also demonstrated effective control and management of Acetamiprid residues in cucumbers, making it a suitable option for pest control in cucumber farming (111). In addition, it has a shorter pre-harvest interval, allowing it to be applied closer to the harvest (112).

On the other hand, this may indicate a greater persistence of this pesticide in cucumbers compared to the other two pesticides, which is consistent with the results of similar studies (108,113). Based on the searches conducted in the present study, there is no specific information on why Acetamiprid may be more persistent in cucumbers. However, the persistence of Acetamiprid in cucumbers could be influenced by several factors, such as its chemical properties, method of application, and interactions with cucumber plants (114). Acetamiprid is highly permeable and can be quickly absorbed and transmitted to all parts of the crop, contributing to its persistence (98). Additionally, the specific metabolic and degradation pathways of this pesticide and its interactions with plant physiology may also play an important role in its persistence (115). However, additional or specialized studies may be needed to fully understand the factors contributing to the persistence of Acetamiprid in plants like cucumber.

Generally, as shown in Figure 3, pesticides can be absorbed by plants through the leaves and roots (116).

The absorption of pesticides by plant roots mainly occurs through the transepithelial absorption process (117). Transepithelial absorption refers to the movement of pesticides through apoplastic and symplastic pathways, in which substances move through the cytoplasm or cell walls of the root's epithelial cells, respectively (118). These pathways play a vital role in the absorption and movement of pesticides within plants, influencing their distribution and potential effects on plant metabolism and growth (119).

In the absorption of pesticides from leaves, simple diffusion has a critical role (120). When a drop of pesticide falls on a leaf, absorption begins at the surface. For the pesticide to reach the cytoplasm, it must move through the cuticle, the cell wall, and finally, the cell membrane. Accordingly, the leaf surface area is a significant factor in the amount of pesticide absorption in plants (121).

To evaluate the potential non-cancer health hazards associated with exposure to pesticides in cucumbers, HQ was calculated (122). An $HQ \leq 1$ suggests that adverse effects are unlikely to occur, indicating a negligible




Figure 3. Absorption pathways of pesticides in cucumber

level of hazard. In contrast, $HQs \geq 1$ indicate a potential for adverse effects, and further toxicological analysis is recommended (123). According to the results reported in Table 5, the HQ values of all three types of pesticides for both children and adults were less than 1 in all countries, implying no appreciable non-cancer health risk (124,125). However, it is important to consider the following points:

- Other exposure routes, such as inhalation and skin contact, may still be present and cannot be disregarded in a non-cancer risk assessment (126,127).
- The per capita consumption of cucumber differs notably across various countries, which can impact human health due to pesticide exposure. High per capita consumption of cucumber increases exposure, and therefore, enhances absorption in the body (128).
- Although all detected pesticides had no potential health risk individually, it should be noted that pesticide exposure through cucumber consumption, due to the cucumber consumption as uncooked foods, can be a serious concern in the long term if they are unwashed (129, 130).
- According to the “cocktail effect” phenomenon, when pesticides are combined in food, they can have a more significant impact on human health than when they are used individually. Therefore, it is essential to conduct comprehensive studies on the combined effects of pesticides rather than focusing solely on the individual effects of each pesticide (131,132).

Furthermore, the results from this study suggest that children face comparatively greater risks than adults. The increased risk for children could be due to their higher

food intake per kilogram of body weight (133), as reported in similar previous studies (134,135).

According to the findings, it is concluded that although the HQ ratio was less than one in all countries, the possible consequences of excess exposure to pesticides deserve further attention and special measures need to be taken to protect children, the most vulnerable population, from pesticide toxicity. The presence of pesticide residues in cucumbers is a critical public health concern that requires ongoing monitoring and innovative detection methods. While current assessments indicate that risks are generally low, continuous vigilance is necessary to mitigate potential health impacts associated with long-term exposure to these chemicals.

One of the most important limitations of this study is the difference in per capita cucumber consumption across different countries, which is influenced by cultural preferences, dietary habits, and economic factors. This variation can impact the calculated risk values related to health and nutrition. Understanding these differences is crucial for health policymakers and researchers when assessing dietary risks and planning interventions tailored to specific populations. It would be more effective to calculate the levels of risk for each country based on its per capita consumption of cucumbers. Also, the amount of agricultural pesticide use varies among countries, with some having implemented strict laws regarding pesticide application in agriculture. This issue should also be taken into consideration when assessing the health risks of pesticides.

Conclusion

In the present report, the pooled concentrations of the investigated chemicals (acetamiprid, chlorpyrifos, and diazinon) in cucumbers were assessed via meta-analysis. The next step involved estimating the corresponding n-CR through the consumption of the vegetable. The analyzed data retrieved from 20 papers indicated pooled concentrations of 140.91, 59.03, and 40.49 µg/kg for acetamiprid, diazinon, and chlorpyrifos, respectively. Depending on some factors such as culture, agricultural methods, pesticide application, economic factors, food security regulations, and politics, the pooled concentrations were obtained differently among the countries. The corresponding health risk calculated using the EPA method for all chemicals indicated an HQ lower than 1, which interprets safe consumption. Although the results indicate safe consumption, considering that cucumber is a fixed product in the diet, its regular consumption may still pose health risks. The implementation of public awareness campaigns, along with legal regulations (e.g., the EU's Farm to Fork strategy) could significantly reduce the health impacts associated with pesticide use. Mitigating the human health impacts of agricultural pesticide use worldwide could be achieved by developing national and international plans for the sustainable use of pesticides, with an emphasis on the gradual elimination of hazardous pesticides, providing low-risk alternatives for high-risk crops, and strengthening periodic training to improve pesticide user performance.

Acknowledgments

We are grateful for the invaluable support provided by the Shahid Beheshti University of Medical Sciences throughout the research process.

Authors' contributions

Conceptualization: Negin Rezaeiashad, Yadolah Fakhri, Mohsen Sadani.

Data curation: Negin Rezaeiashad, Yadolah Fakhri, Mohsen Sadani.

Funding acquisition: Negin Rezaeiashad, Yadolah Fakhri, Mohsen Sadani.

Investigation: Pouria Karimi, Sepideh Sadeghi, Fatemeh Kariminejad, Negin Rezaeiashad, Yadolah Fakhri, Mohsen Sadani.

Methodology: Pouria Karimi, Yadolah Fakhri, Mohsen Sadani, Mohammad Sadeghi, Farshid Soleimani.

Project administration: Pouria Karimi, Sepideh Sadeghi.

Resources: Mohsen Sadani, Mohammad Sadeghi.

Software: Mohsen Sadani, Mohammad Sadeghi.

Supervision: Mohammad Sadeghi.

Validation: Mohsen Sadani, Mohammad Sadeghi.

Visualization: Mohsen Sadani, Mohammad Sadeghi.

Writing—original draft: Pouria Karimi, Sepideh Sadeghi,

Fatemeh Kariminejad, Negin Rezaeiashad, Yadolah Fakhri, Mohsen Sadani, Mohammad Sadeghi.

Writing—review & editing: Pouria Karimi, Sepideh Sadeghi, Fatemeh Kariminejad, Negin Rezaeiashad, Yadolah Fakhri, Mohsen Sadani, Mohammad Sadeghi.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Ethical issues

This study was approved by the Ethics Committee of the Student Research Committee at Shahid Beheshti University of Medical Sciences.

Funding

This project was approved in the Student Research Committee at Shahid Beheshti University of Medical Sciences [1400/66628].

References

1. Alipour V, Mahmoudi I, Borzoei M, Mehri F, Sarkhosh M, Limam I, et al. Concentration of potentially toxic elements (PTEs) in rapid coffee products in Bandar Abbas, Iran: probabilistic non-carcinogenic and carcinogenic risk assessment. *Biol Trace Elem Res*. 2024. doi: [10.1007/s12011-024-04228-y](https://doi.org/10.1007/s12011-024-04228-y).
2. Ualiyeva RM, Kukusheva AN, Insebayeva MK, Akhmetov KK, Zhangazin SB, Krykbayeva MS. Agrotechnological methods of plant feeders applying for spring wheat agrocenoses-North-Eastern Kazakhstan varieties. *J Water Land Dev*. 2022(55):28-40. doi: [10.24425/jwld.2022.142301](https://doi.org/10.24425/jwld.2022.142301).
3. Rezaeigolestani M, Hashemi M. A review of pesticide residues in agricultural and food products of Iran. *J Nutr Fasting Health*. 2018;6(1):1-6. doi: [10.22038/jnfh.2018.33593.1125](https://doi.org/10.22038/jnfh.2018.33593.1125).
4. Ualiyeva RM, Kaverina MM, Ivanko LN, Zhangazin SB. Assessment of spring wheat varieties for pest resistance. *Online J Biol Sci*. 2023;23(4):489-503. doi: [10.3844/ojbsci.2023.489.503](https://doi.org/10.3844/ojbsci.2023.489.503).
5. Alexandratos N, Bruinsma J. *World Agriculture Towards 2030/2050: the 2012 Revision*. Rome: FAO; 2012. doi: [10.22004/ag.econ.288998](https://doi.org/10.22004/ag.econ.288998).
6. van Dijk M, Morley T, Rau ML, Saghai Y. A meta-analysis of projected global food demand and population at risk of hunger for the period 2010-2050. *Nat Food*. 2021;2(7):494-501. doi: [10.1038/s43016-021-00322-9](https://doi.org/10.1038/s43016-021-00322-9).
7. Atamaleki A, Miranzadeh MB, Mostafaei GR, Akbari H, Iranshahi L, Ghanbari F, et al. Effect of coagulation and sonication on the dissolved air flotation (DAF) process for thickening of biological sludge in wastewater treatment. *Environ Health Eng Manag*. 2020;7(1):59-65. doi: [10.34172/ehem.2020.08](https://doi.org/10.34172/ehem.2020.08).
8. Srivastav AL. Chemical fertilizers and pesticides: role in groundwater contamination. In: Prasad MN, ed. *Agrochemicals Detection, Treatment and Remediation*. Butterworth-Heinemann; 2020. p. 143-59. doi: [10.1016/b978-0-08-103017-2.00006-4](https://doi.org/10.1016/b978-0-08-103017-2.00006-4).

9. Yousefi F, Asadikaram G, Karamouzian S, Abolhassani M, Moazed V, Nematollahi MH. MGMT methylation alterations in brain cancer following organochlorine pesticides exposure. *Environ Health Eng Manag.* 2021;8(1):47-53. doi: [10.34172/ehem.2021.07](https://doi.org/10.34172/ehem.2021.07).
10. Hammood MA, Chamani A, Sobhanardakani S. Health risk assessment of heavy metal exposure from indoor dust: a case study of residential buildings in Isfahan, Iran. *Environ Health Eng Manag.* 2024;11(1):39-50. doi: [10.34172/ehem.2024.06](https://doi.org/10.34172/ehem.2024.06).
11. Chung SW. How effective are common household preparations on removing pesticide residues from fruit and vegetables? A review. *J Sci Food Agric.* 2018;98(8):2857-70. doi: [10.1002/jsfa.8821](https://doi.org/10.1002/jsfa.8821).
12. Amjad A, Saeed MK, Amjad MR, Hameed A, Nawaz H, Ullah N, et al. Effect of ozone, hydrogen peroxide, and chlorine solution in reduction of chlorpyrifos and cypermethrin residues from cauliflower. *Ital J Food Sci.* 2024;36(4):142-50. doi: [10.15586/ijfs.v36i4.2621](https://doi.org/10.15586/ijfs.v36i4.2621).
13. Karami-Mohajeri S, Ahmadipour A, Rahimi HR, Abdollahi M. Adverse effects of organophosphorus pesticides on the liver: a brief summary of four decades of research. *Arh Hig Rada Toksikol.* 2017;68(4):261-75. doi: [10.1515/aiht-2017-68-2989](https://doi.org/10.1515/aiht-2017-68-2989).
14. Amrollahi H, Pazoki R, Imani S. Pesticide multiresidue analysis in tomato and cucumber samples collected from fruit and vegetable markets in Tehran, Iran. *Middle East J Rehabil Health Stud.* 2019;6(1):e64271. doi: [10.5812/mejrh.64271](https://doi.org/10.5812/mejrh.64271).
15. Bahardoust M, Zolfaghari Dehkharghani M, Ebrahimi P, Najafirashed M, Mousavi S, Haghmoradi M, et al. Effect of ABO blood group on postoperative overall survival and recurrence-free survival rate in patients with hepatocellular carcinoma after hepatectomy: a multi-center retrospective cohort study. *BMC Surg.* 2023;23(1):324. doi: [10.1186/s12893-023-02236-8](https://doi.org/10.1186/s12893-023-02236-8).
16. Leonel Javeres MN, Habib R, Judith Laure N, Abbas Shah ST, Valis M, Kuca K, et al. Chronic exposure to organophosphates pesticides and risk of metabolic disorder in cohort from Pakistan and Cameroon. *Int J Environ Res Public Health.* 2021;18(5):2310. doi: [10.3390/ijerph18052310](https://doi.org/10.3390/ijerph18052310).
17. Oostingh GJ, Wichmann G, Schmittner M, Lehmann I, Duschl A. The cytotoxic effects of the organophosphates chlorpyrifos and diazinon differ from their immunomodulating effects. *J Immunotoxicol.* 2009;6(2):136-45. doi: [10.1080/15476910902977407](https://doi.org/10.1080/15476910902977407).
18. Mansour SA, Belal MH, Abou-Arab AA, Gad MF. Monitoring of pesticides and heavy metals in cucumber fruits produced from different farming systems. *Chemosphere.* 2009;75(5):601-9. doi: [10.1016/j.chemosphere.2009.01.058](https://doi.org/10.1016/j.chemosphere.2009.01.058).
19. Onyeaka H, Anumudu CK, Okolo CA, Anyogu A, Odeyemi O, Bassey AP. A review of the top 100 most cited papers on food safety. *Qual Assur Saf Crop Foods.* 2022;14(4):91-104. doi: [10.15586/qas.v14i4.1124](https://doi.org/10.15586/qas.v14i4.1124).
20. Kashi G, Nourieh N, Mostashari P, Khushab F. Optimization of extraction conditions and determination of the chlorpyrifos, diazinon, and malathion residues in environment samples: fruit (apple, orange, and tomato). *Food Chem X.* 2021;12:100163. doi: [10.1016/j.fochx.2021.100163](https://doi.org/10.1016/j.fochx.2021.100163).
21. Sidhu GK, Singh S, Kumar V, Dhanjal DS, Datta S, Singh J. Toxicity, monitoring and biodegradation of organophosphate pesticides: a review. *Crit Rev Environ Sci Technol.* 2019;49(13):1135-87. doi: [10.1080/10643389.2019.1565554](https://doi.org/10.1080/10643389.2019.1565554).
22. Hu L, Wang X, Zou Y, Wu D, Gao G, Zhong Z, et al. Effects of inorganic and organic selenium intervention on resistance of radish to arsenic stress. *Ital J Food Sci.* 2022;34(1):44-58. doi: [10.15586/ijfs.v34i1.2105](https://doi.org/10.15586/ijfs.v34i1.2105).
23. Siripongvutikorn S, Usawakesmanee W, Pisuchpen S, Khatcharin N, Rujirapong C. Quality changes during storage in Thai indigenous leafy vegetable, Liang leaves (*Gnetum gnemon* var. *tenerum*) after different preparation methods. *Ital J Food Sci.* 2023;35(3):1-16. doi: [10.15586/ijfs.v35i3.2346](https://doi.org/10.15586/ijfs.v35i3.2346).
24. Witczak A, Pohoryło A, Abdel-Gawad H, Cybulski J. Residues of some organophosphorus pesticides on and in fruits and vegetables available in Poland, an assessment based on the European Union regulations and health assessment for human populations. *Phosphorus Sulfur Silicon Relat Elem.* 2018;193(11):711-20. doi: [10.1080/10426507.2018.1492921](https://doi.org/10.1080/10426507.2018.1492921).
25. Omwenga I, Kanja L, Zomer P, Louisse J, Rietjens I, Mol H. Organophosphate and carbamate pesticide residues and accompanying risks in commonly consumed vegetables in Kenya. *Food Addit Contam Part B Surveill.* 2021;14(1):48-58. doi: [10.1080/19393210.2020.1861661](https://doi.org/10.1080/19393210.2020.1861661).
26. Kaur M, Sharma P. Recent advances in cucumber (*Cucumis sativus* L.). *J Hortic Sci Biotechnol.* 2022;97(1):3-23. doi: [10.1080/14620316.2021.1945956](https://doi.org/10.1080/14620316.2021.1945956).
27. Mallick PK. Evaluating potential importance of cucumber (*Cucumis sativus* L.-Cucurbitaceae): a brief review. *Int J Appl Sci Biotechnol.* 2022;10(1):12-5. doi: [10.3126/ijasbt.v10i1.44152](https://doi.org/10.3126/ijasbt.v10i1.44152).
28. Ahmed GA. Controlling of *Fusarium* wilt of cucumber by antagonistic bacteria. *J Life Sci.* 2010;4(7):16-21.
29. Mohamed AO, Mater AA, Hammad AM, Ishag AE, El Tayeb EM, Dahab AA. Pesticide residues detected on tomato and cucumber fruits grown in greenhouse farms in Khartoum state, Sudan. *Int J Life Sci Res.* 2018;6(3):472-81.
30. Ostovan H, Yazdanpak A, Hesami S, Gheibi M. Assessment of four pesticide residues (diazinon, imidacloprid, pirimicarb and acetamiprid) in cucumber under greenhouse condition of Iran (Fars province). *IAU Entomological Research Journal.* 2018;10(2):139-48.
31. LingSJ, MahatNA. Presence of Selected Organophosphorous Pesticide Residues in Raw Cucumber and Tomato Samples in Taman Universiti, Johor [dissertation]. Universiti Teknologi Malaysia; 2014. p. 32-9.
32. Fakhri Y, Atamaleki A, Asadi A, Ghasemi SM, Mousavi Khaneghah A. Bioaccumulation of potentially toxic elements (PTEs) in muscle *Tilapia* spp fish: a systematic review, meta-analysis, and non-carcinogenic risk assessment. *Toxin Rev.* 2021;40(4):473-83. doi: [10.1080/15569543.2019.1690518](https://doi.org/10.1080/15569543.2019.1690518).
33. Atamaleki A, Yazdanbakhsh A, Fakhri Y, Salem A, Ghorbanian M, Mousavi Khaneghah A. A systematic review and meta-analysis to investigate the correlation vegetable irrigation with wastewater and concentration of potentially toxic elements (PTEs): a case study of Spinach (*Spinacia oleracea*) and radish (*Raphanus raphanistrum* subsp. *sativus*). *Biol Trace Elem Res.* 2021;199(2):792-9. doi: [10.1007/s12011-020-02181-0](https://doi.org/10.1007/s12011-020-02181-0).
34. Zamanpour S, Noori SM, Shokrollahi Yancheshmeh B,

- Afshari A, Hashemi M. A systematic review to introduce the most effective postbiotics derived from probiotics for aflatoxin detoxification in vitro. *Ital J Food Sci.* 2023;35(4):31-49. doi: [10.15586/ijfs.v35i4.2369](https://doi.org/10.15586/ijfs.v35i4.2369).
35. Ali S, Mehri F, Nasiri R, Limam I, Fakhri Y. Fluoride in raw rice (*Oryza sativa*): a global systematic review and probabilistic health risk assessment. *Biol Trace Elem Res.* 2024;202(9):4324-33. doi: [10.1007/s12011-023-04004-4](https://doi.org/10.1007/s12011-023-04004-4).
 36. Atamaleki A, Yazdanbakhsh A, Gholizadeh A, Naimi N, Karimi P, Thai VN, et al. Concentration of potentially harmful elements (PHEs) in eggplant vegetable (*Solanum melongena*) irrigated with wastewater: a systematic review and meta-analysis and probabilistic health risk assessment. *Int J Environ Health Res.* 2022;32(7):1419-31. doi: [10.1080/09603123.2021.1887461](https://doi.org/10.1080/09603123.2021.1887461).
 37. Karimi P, Yazdanbakhsh A, Atamaleki A, Kariminejad F, Ahmadian M, Fallah S. Global evaluation of potentially harmful elements (PHEs) in potato and carrot irrigated by wastewater: a systematic review, meta-analysis, and health risk assessment. *Environ Health Eng Manag.* 2021;8(4):245-55. doi: [10.34172/ehem.2021.28](https://doi.org/10.34172/ehem.2021.28).
 38. Food and Agriculture Organization (FAO). Pesticides Use [Internet]. FAO; 2020. Available from: <https://www.fao.org/faostat/en/#data/RP>.
 39. Regional Screening Levels (RSLs) [Internet]. EPA. 2020. Available from: <https://www.epa.gov/risk/regional-screening-levels-rsls-generic-tables>.
 40. Fu D, Zhang S, Wang M, Liang X, Xie Y, Zhang Y, et al. Dissipation behavior, residue distribution and dietary risk assessment of cyromazine, acetamiprid and their mixture in cowpea and cowpea field soil. *J Sci Food Agric.* 2020;100(12):4540-8. doi: [10.1002/jsfa.10495](https://doi.org/10.1002/jsfa.10495).
 41. Naimi N, Pilevar Z, Ranaei V, Mahmudiono T, Fakhri Y, Paseban A, et al. The concentration of potentially toxic elements (PTEs) in apple fruit: a global systematic review, meta-analysis, and health risk assessment. *Environ Sci Pollut Res Int.* 2022;29(36):54013-24. doi: [10.1007/s11356-022-21158-1](https://doi.org/10.1007/s11356-022-21158-1).
 42. Król S, Zabiegała B, Namieśnik J. Monitoring VOCs in atmospheric air I. On-line gas analyzers. *Trends Analyt Chem.* 2010;29(9):1092-100. doi: [10.1016/j.trac.2010.05.007](https://doi.org/10.1016/j.trac.2010.05.007).
 43. Sharma A, Kumar V, Shahzad B, Tanveer M, Sidhu GP, Handa N, et al. Worldwide pesticide usage and its impacts on ecosystem. *SN Appl Sci.* 2019;1(11):1446. doi: [10.1007/s42452-019-1485-1](https://doi.org/10.1007/s42452-019-1485-1).
 44. Liu EM, Huang J. Risk preferences and pesticide use by cotton farmers in China. *J Dev Econ.* 2013;103:202-15. doi: [10.1016/j.jdeveco.2012.12.005](https://doi.org/10.1016/j.jdeveco.2012.12.005).
 45. Yu R, Liu Q, Liu J, Wang Q, Wang Y. Concentrations of organophosphorus pesticides in fresh vegetables and related human health risk assessment in Changchun, Northeast China. *Food Control.* 2016;60:353-60. doi: [10.1016/j.foodcont.2015.08.013](https://doi.org/10.1016/j.foodcont.2015.08.013).
 46. Vlassi E, Bempelou E, Liapis K, Arapis G. Consumer safety evaluation after monitoring of endocrine disruptor pesticide residues: a case study of Thessaly, Central Greece. *Toxicol Environ Chem.* 2020;102(1-4):105-23. doi: [10.1080/02772248.2020.1770256](https://doi.org/10.1080/02772248.2020.1770256).
 47. Tarawneh IN, Alawi MA, Sapah RH, Abu Shmeis RM. Pesticide residues in commonly consumed fruits and vegetables in Jordan and their associated health risk assessments. *Jordan J Chem.* 2019;14(2):69-80.
 48. Fatunsin OT, Oyeyiola AO, Moshood MO, Akanbi LM, Fadahunsi DE. Dietary risk assessment of organophosphate and carbamate pesticide residues in commonly eaten food crops. *Sci Afr.* 2020;8:e00442. doi: [10.1016/j.sciaf.2020.e00442](https://doi.org/10.1016/j.sciaf.2020.e00442).
 49. Shokrzadeh M, Enayati A, Mokarami H, Ziar A. Assessment of diazinon and malathion residues in cucumber produced in Mazandaran, Iran. *J Mazandaran Univ Med Sci.* 2017;27(152):143-52. [Persian].
 50. Crépet A, Luong TM, Baines J, Boon PE, Ennis J, Kennedy M, et al. An international probabilistic risk assessment of acute dietary exposure to pesticide residues in relation to codex maximum residue limits for pesticides in food. *Food Control.* 2021;121:107563. doi: [10.1016/j.foodcont.2020.107563](https://doi.org/10.1016/j.foodcont.2020.107563).
 51. Nasiri A, Amirahmadi M, Mousavi Z, Shoeibi S, Khajeamiri A, Kobarfard F. A multi residue GC-MS method for determination of 12 pesticides in cucumber. *Iran J Pharm Res.* 2016;15(4):809-16.
 52. Mosleh YY, Mofeed J, Almagrabi OA, Mousa T, Kadasa NM. Dietary intake of pesticides based on vegetable consumption: a case study, Jeddah, Kingdom of Saudi Arabia. *Life Sci J.* 2014;11(7):680-8.
 53. Lozowicka B, Abzeitova E, Sagitov A, Kaczynski P, Toleubayev K, Li A. Studies of pesticide residues in tomatoes and cucumbers from Kazakhstan and the associated health risks. *Environ Monit Assess.* 2015;187(10):609. doi: [10.1007/s10661-015-4818-6](https://doi.org/10.1007/s10661-015-4818-6).
 54. Jamshidi B, Mohajerani E, Jamshidi J. Developing a Vis/NIR spectroscopic system for fast and non-destructive pesticide residue monitoring in agricultural product. *Measurement.* 2016;89:1-6. doi: [10.1016/j.measurement.2016.03.069](https://doi.org/10.1016/j.measurement.2016.03.069).
 55. Ganjezadeh Rohani F, Mahdavi V, Aminaei MM. Investigation on diazinon and oxydemeton-methyl residues in cucumbers grown in Kerman greenhouses. *Environ Monit Assess.* 2014;186(7):3995-9. doi: [10.1007/s10661-014-3674-0](https://doi.org/10.1007/s10661-014-3674-0).
 56. El-Saeid MH, Selim MT. Multiresidue analysis of 86 pesticides using gas chromatography mass spectrometry: II-nonleafy vegetables. *J Chem.* 2013;2013(1):727149. doi: [10.1155/2013/727149](https://doi.org/10.1155/2013/727149).
 57. Davoodi D, Hassanzadeh-Khayyat M, Asgharian Rezaei M, Mohajer SA. Preparation, evaluation and application of diazinon imprinted polymers as the sorbent in molecularly imprinted solid-phase extraction and liquid chromatography analysis in cucumber and aqueous samples. *Food Chem.* 2014;158:421-8. doi: [10.1016/j.foodchem.2014.02.144](https://doi.org/10.1016/j.foodchem.2014.02.144).
 58. Darvishnejad F, Raofi JB, Ghani M. MIL-101 (Cr) @ graphene oxide-reinforced hollow fiber solid-phase microextraction coupled with high-performance liquid chromatography to determine diazinon and chlorpyrifos in tomato, cucumber and agricultural water. *Anal Chim Acta.* 2020;1140:99-110. doi: [10.1016/j.aca.2020.10.015](https://doi.org/10.1016/j.aca.2020.10.015).
 59. Bazmandegan-Shamili A, Haji Shabani AM, Dadfarnia S, Rohani Moghadam M, Saeidi M. Preparation of magnetic mesoporous silica composite for the solid-phase microextraction of diazinon and malathion before their determination by high-performance liquid chromatography. *J Sep Sci.* 2017;40(8):1731-8. doi: [10.1002/jssc.201601339](https://doi.org/10.1002/jssc.201601339).
 60. Bakırcı GT, Yaman Acay DB, Bakırcı F, Ötleş S. Pesticide residues in fruits and vegetables from the Aegean region,

- Turkey. Food Chem. 2014;160:379-92. doi: [10.1016/j.foodchem.2014.02.051](https://doi.org/10.1016/j.foodchem.2014.02.051).
61. Bagheri H, Es'haghi A, Es-Haghi A, Mesbahi N. A high-throughput approach for the determination of pesticide residues in cucumber samples using solid-phase microextraction on 96-well plate. Anal Chim Acta. 2012;740:36-42. doi: [10.1016/j.aca.2012.06.001](https://doi.org/10.1016/j.aca.2012.06.001).
 62. Galani YJ, Houbraken M, Wumbei A, Djeugap JF, Fotio D, Gong YY, et al. Monitoring and dietary risk assessment of 81 pesticide residues in 11 local agricultural products from the 3 largest cities of Cameroon. Food Control. 2020;118:107416. doi: [10.1016/j.foodcont.2020.107416](https://doi.org/10.1016/j.foodcont.2020.107416).
 63. Quijano L, Yusà V, Font G, Pardo O. Chronic cumulative risk assessment of the exposure to organophosphorus, carbamate and pyrethroid and pyrethrin pesticides through fruit and vegetables consumption in the region of Valencia (Spain). Food Chem Toxicol. 2016;89:39-46. doi: [10.1016/j.fct.2016.01.004](https://doi.org/10.1016/j.fct.2016.01.004).
 64. Siddique Z, Malik AU, Asi MR, Inam-Ur-Raheem M, Iqbal M, Abdullah M. Impact of sonolytic ozonation (O₃/US) on degradation of pesticide residues in fresh vegetables and fruits: case study of Faisalabad, Pakistan. Ultrason Sonochem. 2021;79:105799. doi: [10.1016/j.ultsonch.2021.105799](https://doi.org/10.1016/j.ultsonch.2021.105799).
 65. Hu Z. What socio-economic and political factors lead to global pesticide dependence? A critical review from a social science perspective. Int J Environ Res Public Health. 2020;17(21):8119. doi: [10.3390/ijerph17218119](https://doi.org/10.3390/ijerph17218119).
 66. Sarkar S, Gil JD, Keeley J, Jansen K. The Use of Pesticides in Developing Countries and Their Impact on Health and the Right to Food. European Union; 2021.
 67. Larson SJ. Pesticides in Surface Waters: Distribution, Trends, and Governing Factors. CRC Press; 2019. doi: [10.1201/9780429062797](https://doi.org/10.1201/9780429062797).
 68. Grafton-Cardwell EE, Leonard JT, Daugherty MP, Headrick DH. Mating disruption of the California red scale, *Aonidiella aurantii* (Hemiptera: Diaspididae) in central California citrus. J Econ Entomol. 2021;114(6):2421-9. doi: [10.1093/jee/toab188](https://doi.org/10.1093/jee/toab188).
 69. Deacon S, Norman S, Nicolette J, Reub G, Greene G, Osborn R, et al. Integrating ecosystem services into risk management decisions: case study with Spanish citrus and the insecticide chlorpyrifos. Sci Total Environ. 2015;505:732-9. doi: [10.1016/j.scitotenv.2014.10.034](https://doi.org/10.1016/j.scitotenv.2014.10.034).
 70. Thuy TN, Thanh Phong L, van Kien N, Phuong Dong LT. The impacts of paraquat and chlorpyrifos in agricultural production on environment and farmers' health in a Giang province, Vietnam. Imp J Interdiscip Res. 2016;2(6):860-6.
 71. Garcerá C, Moltó E, Chueca P. Effect of spray volume of two organophosphate pesticides on coverage and on mortality of California red scale *Aonidiella aurantii* (Maskell). Crop Prot. 2011;30(6):693-7. doi: [10.1016/j.cropro.2011.02.019](https://doi.org/10.1016/j.cropro.2011.02.019).
 72. Ahmed MT, Greish S, Ismail SM, Mosleh Y, Loutfy NM, El Doussouki A. Dietary intake of pesticides based on vegetable consumption in Ismailia, Egypt: a case study. Hum Ecol Risk Assess. 2014;20(3):779-88. doi: [10.1080/10807039.2013.775893](https://doi.org/10.1080/10807039.2013.775893).
 73. Eissa F, Al-Sisi M, Ghanem K. Occurrence and ecotoxicological risk assessment of pesticides in sediments of the Rosetta branch, Nile River, Egypt. J Environ Sci (China). 2022;118:21-31. doi: [10.1016/j.jes.2021.08.047](https://doi.org/10.1016/j.jes.2021.08.047).
 74. Salem SH, Fatah AE, Abdel-Rahman GN, Fouzy Ū, Marrez D. Screening for pesticide residues in soil and crop samples in Egypt. Egypt J Chem. 2021;64(5):2525-32. doi: [10.21608/ejchem.2021.64117.3374](https://doi.org/10.21608/ejchem.2021.64117.3374).
 75. Ibrahim EA, Shalaby SEM. Screening and assessing of pesticide residues and their health risks in vegetable field soils from the eastern Nile Delta, Egypt. Toxicol Rep. 2022;9:1281-90. doi: [10.1016/j.toxrep.2022.06.004](https://doi.org/10.1016/j.toxrep.2022.06.004).
 76. Abuzaid AS, Abdel-Salam MA, Ahmad AF, Fathy HA, Fadl ME, Scopa A. Effect of marginal-quality irrigation on accumulation of some heavy metals (Mn, Pb, and Zn) in Typic Torripsamment soils and food crops. Sustainability. 2022;14(3):1067. doi: [10.3390/su14031067](https://doi.org/10.3390/su14031067).
 77. Hathout A, Amer M, Hussain O, Yassen AA, Mossa AT, Elgohary MR. Estimation of the most widespread pesticides in agricultural soils collected from some Egyptian governorates. Egypt J Chem. 2022;65(1):35-44. doi: [10.21608/ejchem.2021.72087.3591](https://doi.org/10.21608/ejchem.2021.72087.3591).
 78. Bakr A, Mahmoud HA, Ghanem KM, Eissa FI. Monitoring and risk assessment of organophosphorus pesticide residues in surface and drinking water in some Egyptian governorates. Egypt J Chem. 2023;66(13):77-87. doi: [10.21608/ejchem.2023.210533.7967](https://doi.org/10.21608/ejchem.2023.210533.7967).
 79. Eissa FI, Mahmoud HA, Massoud ON, Ghanem KM, Gomaa IM. Biodegradation of chlorpyrifos by microbial strains isolated from agricultural wastewater. J Am Sci. 2014;10(3):98-108.
 80. Crane AL, Abdel Rasoul G, Ismail AA, Hendy O, Bonner MR, Lasarev MR, et al. Longitudinal assessment of chlorpyrifos exposure and effect biomarkers in adolescent Egyptian agricultural workers. J Expo Sci Environ Epidemiol. 2013;23(4):356-62. doi: [10.1038/jes.2012.113](https://doi.org/10.1038/jes.2012.113).
 81. Ibrahim A, Khorshed MA, Eshmawy MR, Kandil MA. Pesticide residues in horticultural crops from domestic markets of Giza, Egypt: occurrence and risk assessment. Plant Arch. 2020;20(1):3862-9.
 82. Wu X, Li J, Zhou Z, Lin Z, Pang S, Bhatt P, et al. Environmental occurrence, toxicity concerns, and degradation of diazinon using a microbial system. Front Microbiol. 2021;12:717286. doi: [10.3389/fmicb.2021.717286](https://doi.org/10.3389/fmicb.2021.717286).
 83. Dahmardeh Behrooz R, Esmaili-Sari A, Urbaniak M, Chakraborty P. Assessing diazinon pollution in the three major rivers flowing into the Caspian Sea (Iran). Water. 2021;13(3):335. doi: [10.3390/w13030335](https://doi.org/10.3390/w13030335).
 84. Akbarlou Z, Alipour V, Heidari M, Dindarloo K. Adsorption of diazinon from aqueous solutions onto an activated carbon sample produced in Iran. Environ Health Eng Manag. 2017;4(2):93-9. doi: [10.15171/ehem.2017.13](https://doi.org/10.15171/ehem.2017.13).
 85. Kermani M, Dowlati M, Gholami M, Sobhi HR, Azari A, Esrafil A, et al. A global systematic review, meta-analysis and health risk assessment on the quantity of malathion, diazinon and chlorpyrifos in vegetables. Chemosphere. 2021;270:129382. doi: [10.1016/j.chemosphere.2020.129382](https://doi.org/10.1016/j.chemosphere.2020.129382).
 86. Morteza Z, Mousavi SB, Baghestani MA, Aitio A. An assessment of agricultural pesticide use in Iran, 2012-2014. J Environ Health Sci Eng. 2017;15:10. doi: [10.1186/s40201-017-0272-4](https://doi.org/10.1186/s40201-017-0272-4).
 87. Nasrollahi M, Pourbabaei AA, Etesami H, Talebi K. Diazinon degradation by bacterial endophytes in rice plant (*Oryza sativa* L.): a possible reason for reducing the efficiency of diazinon in the control of the rice stem-borer. Chemosphere. 2020;246:125759. doi: [10.1016/j.chemosphere.2019.125759](https://doi.org/10.1016/j.chemosphere.2019.125759).
 88. Ebrahimzadeh G, Alimohammadi M, Rezaei Kahkha MR, Mahvi AH. Contamination level and human non-

- carcinogenic risk assessment of diazinon pesticide residue in drinking water resources—a case study, Iran. *Int J Environ Anal Chem.* 2022;102(16):4726-37. doi: [10.1080/03067319.2020.1789609](https://doi.org/10.1080/03067319.2020.1789609).
89. Dehghani R, Shayeghi M, Eslami H, Moosavi SG, Rabbani D, Hossein Shahi D. Determination of organophosphorus pesticides (diazinon and chlorpyrifos) in water resources in Barzok, Kashan. *Zahedan J Res Med Sci.* 2012;14(10):e93196.
 90. Abadi B. The determinants of cucumber farmers' pesticide use behavior in central Iran: Implications for the pesticide use management. *J Clean Prod.* 2018;205:1069-81. doi: [10.1016/j.jclepro.2018.09.147](https://doi.org/10.1016/j.jclepro.2018.09.147).
 91. Saggu S, Rehman H, Alzeiber FM, Aziz A. Current situation of pesticide consumption and poisoning in Saudi Arabia. *J Entomol Zool Stud.* 2016;4(3):153-8.
 92. Alokail MS, Abd-Alrahman SH, Alnaami AM, Hussain SD, Amer OE, Elhalwagy ME, et al. Regional variations in pesticide residue detection rates and concentrations in Saudi Arabian crops. *Toxics.* 2023;11(9):798. doi: [10.3390/toxics11090798](https://doi.org/10.3390/toxics11090798).
 93. Al-Wabel M, El-Saeid MH, El-Naggar AH, Al-Romian FA, Osman K, Elnazi K, et al. Spatial distribution of pesticide residues in the groundwater of a condensed agricultural area. *Arab J Geosci.* 2016;9(2):120. doi: [10.1007/s12517-015-2122-y](https://doi.org/10.1007/s12517-015-2122-y).
 94. Ramadan MF, Abdel-Hamid MM, Altorgoman MM, AlGaramah HA, Alawi MA, Shati AA, et al. Evaluation of pesticide residues in vegetables from the Asir region, Saudi Arabia. *Molecules.* 2020;25(1):205. doi: [10.3390/molecules25010205](https://doi.org/10.3390/molecules25010205).
 95. Asiry KA, Huda MN, Mousa MA. Abundance and population dynamics of the key insect pests and agronomic traits of tomato (*Solanum lycopersicon* L.) varieties under different planting densities as a sustainable pest control method. *Horticulturae.* 2022;8(10):976. doi: [10.3390/horticulturae8100976](https://doi.org/10.3390/horticulturae8100976).
 96. Azeem MI, Alhafi Alotaibi B. Farmers' beliefs and concerns about climate change, and their adaptation behavior to combat climate change in Saudi Arabia. *PLoS One.* 2023;18(1):e0280838. doi: [10.1371/journal.pone.0280838](https://doi.org/10.1371/journal.pone.0280838).
 97. Hajjar MJ. Obsolete pesticides in Saudi Arabia: problems, prevention and disposal. *MOJ Toxicol.* 2015;1(2):00009. doi: [10.15406/mojt.2015.01.00009](https://doi.org/10.15406/mojt.2015.01.00009).
 98. Zuščíková L, Bažány D, Greifová H, Knížatová N, Kováčik A, Lukáč N, et al. Screening of toxic effects of neonicotinoid insecticides with a focus on acetamiprid: a review. *Toxics.* 2023;11(7):598. doi: [10.3390/toxics11070598](https://doi.org/10.3390/toxics11070598).
 99. Geng C, Watson GB, Sparks TC. Nicotinic acetylcholine receptors as spinosyn targets for insect pest management. *Adv In Insect Phys.* 2013;44:101-210. doi: [10.1016/b978-0-12-394389-7.00003-x](https://doi.org/10.1016/b978-0-12-394389-7.00003-x).
 100. Sahin C, Karpuzcu ME. Mitigation of organophosphate pesticide pollution in agricultural watersheds. *Sci Total Environ.* 2020;710:136261. doi: [10.1016/j.scitotenv.2019.136261](https://doi.org/10.1016/j.scitotenv.2019.136261).
 101. Erdogan C, Ozdem A, Alpken YN, Demiroz D. Resistance to common insecticides and mechanisms of resistance in *Aphis pomi* De Geer (Hemiptera: Aphididae), in apple orchards in Turkey. *Phytoparasitica.* 2023;51(2):323-35. doi: [10.1007/s12600-022-01042-x](https://doi.org/10.1007/s12600-022-01042-x).
 102. Balkan T, Yılmaz Ö. Investigation of insecticide residues in potato grown in Türkiye by LC-MS/MS and GC-MS and health risk assessment. *Turk J Entomol.* 2022;46(4):481-500. doi: [10.16970/entotod.1201475](https://doi.org/10.16970/entotod.1201475).
 103. Sungur Ş, Tunur Ç. Investigation of pesticide residues in vegetables and fruits grown in various regions of Hatay, Turkey. *Food Addit Contam Part B Surveill.* 2012;5(4):265-7. doi: [10.1080/19393210.2012.704597](https://doi.org/10.1080/19393210.2012.704597).
 104. Toptancı İ, Kiralan M, Ramadan MF. Levels of pesticide residues in fruits and vegetables in the Turkish domestic markets. *Environ Sci Pollut Res.* 2021;28(29):39451-7. doi: [10.1007/s11356-021-13538-w](https://doi.org/10.1007/s11356-021-13538-w).
 105. Kaçikoç M, Censur M. Environmental monitoring of pesticide residues in surface waters of Büyük Menderes River. *Sigma J Eng Nat Sc.* 2022;40(4):762-71. doi: [10.14744/sigma.2022.00092](https://doi.org/10.14744/sigma.2022.00092).
 106. Abu-Duka AB, Mohammadali MT. Study of the effectiveness of pesticides thiamethoxam and acetamiprid against cabbage aphid *Brevicoryne brassicae* and measure the residue of acetamiprid in leaves and soil of cabbage using HPLC. *Plant Cell Biotechnol Mol Biol.* 2021;22(19-20):123-9.
 107. Derafshi M, Black C, Agnew R. Pesticide knowledge, attitudes, and clothing practices of Turkish farmers. *Environmental Management and Sustainable Development.* 2017;6(1):149-67. doi: [10.5296/emsd.v6i1.10271](https://doi.org/10.5296/emsd.v6i1.10271).
 108. Ahmed MA, Hashem DA, Ahmed SA, Khalid NS. Monitoring of pesticide residues in cucumber samples marketed in Egypt. *J Plant Prot Pathol.* 2019;10(4):225-8.
 109. Rubesh GS, Kalyanasundaram A, Ambethgar V, Kuppusamy S, Mohanasundaram A. Pesticide usage patterns and farmer awareness among Solanaceae crop farmers in Tiruchirappalli. *Pharma Innov.* 2023;12(7):2675-80.
 110. Noreen S, Khan IM, Khan MS, Zarnaab B, Gul I, Khan MZ, et al. Comparative valuation of the chlorpyrifos, acetamiprid, and lambda-cyhalothrin toxicity and their hematological and histopathological consequences in pigeons. *Environ Sci Pollut Res Int.* 2023;30(40):92817-29. doi: [10.1007/s11356-023-28515-8](https://doi.org/10.1007/s11356-023-28515-8).
 111. Salehi Mishani H, Jalalizand A, Modaresi M. Investigating the effect of dichlorvos and acetamiprid residues in greenhouse cucumber on biochemical parameters and protective role of colostrum. *J Res Med Sci.* 2023;28:52. doi: [10.4103/jrms.jrms_2_23](https://doi.org/10.4103/jrms.jrms_2_23).
 112. Lazić SA, Šunjka DR, Jovanov PA, Vuković SL, Guzsvány V. LC-MS/MS determination of acetamiprid residues in sweet cherries. *Rom Biotechnol Lett.* 2018;23(1):13317-26.
 113. AlSaikhan WH, Almatroodi SA, Almatroudi A, Alsahli MA, Rahmani AH. Pesticide residue measurement in commonly used vegetables using the QuEChERS method. *Pharmacognosy Journal.* 2021;13(1):142-9. doi: [10.5530/pj.2021.13.20](https://doi.org/10.5530/pj.2021.13.20).
 114. Wang X, Yan M, Zhou J, Song W, Xiao Y, Cui C, et al. Delivery of acetamiprid to tea leaves enabled by porous silica nanoparticles: efficiency, distribution and metabolism of acetamiprid in tea plants. *BMC Plant Biol.* 2021;21(1):337. doi: [10.1186/s12870-021-03120-4](https://doi.org/10.1186/s12870-021-03120-4).
 115. Zhang A, Xu L, Liu Z, Zhang J, Han L, Zhao K. The effects of acetamiprid multigeneration stress on metabolism and physiology of *Aphis glycines* Matsumura (Hemiptera: Aphididae). *Arch Insect Biochem Physiol.* 2022;110(4):e21903. doi: [10.1002/arch.21903](https://doi.org/10.1002/arch.21903).
 116. Zhang C, Yao FE, Liu YW, Chang HQ, Li ZJ, Xue JM. Uptake and translocation of organic pollutants in plants: a review. *J Integr Agric.* 2017;16(8):1659-68. doi: [10.1016/](https://doi.org/10.1016/)

- s2095-3119(16)61590-3.
117. Denecke S, Swevers L, Douris V, Vontas J. How do oral insecticidal compounds cross the insect midgut epithelium? *Insect Biochem Mol Biol.* 2018;103:22-35. doi: [10.1016/j.ibmb.2018.10.005](https://doi.org/10.1016/j.ibmb.2018.10.005).
 118. Nguyen TN, Pham CV, Chowdhury R, Patel S, Jaysawal SK, Hou Y, et al. Development of blueberry-derived extracellular nanovesicles for immunomodulatory therapy. *Pharmaceutics.* 2023;15(8):2115. doi: [10.3390/pharmaceutics15082115](https://doi.org/10.3390/pharmaceutics15082115).
 119. Barberon M, Geldner N. Radial transport of nutrients: the plant root as a polarized epithelium. *Plant Physiol.* 2014;166(2):528-37. doi: [10.1104/pp.114.246124](https://doi.org/10.1104/pp.114.246124).
 120. Wang CJ, Liu ZQ. Foliar uptake of pesticides—present status and future challenge. *Pestic Biochem Physiol.* 2007;87(1):1-8. doi: [10.1016/j.pestbp.2006.04.004](https://doi.org/10.1016/j.pestbp.2006.04.004).
 121. Liao P, Maoz I, Shih ML, Lee JH, Huang XQ, Morgan JA, et al. Emission of floral volatiles is facilitated by cell-wall non-specific lipid transfer proteins. *Nat Commun.* 2023;14(1):330. doi: [10.1038/s41467-023-36027-9](https://doi.org/10.1038/s41467-023-36027-9).
 122. US Environmental Protection Agency (EPA). Regional Screening Table - User's Guide, Mid-Atlantic Risk Assessment. 2019. Available from: <https://www.epa.gov/risk/regional-screening-levels-rsls-generic-tables>.
 123. Tokatli C, Ustaoglu F. Health risk assessment of toxicants in Meriç river delta wetland, Thrace region, Turkey. *Environ Earth Sci.* 2020;79(18):426. doi: [10.1007/s12665-020-09171-4](https://doi.org/10.1007/s12665-020-09171-4).
 124. Kukusamude C, Sricharoen P, Limchoowong N, Kongsri S. Heavy metals and probabilistic risk assessment via rice consumption in Thailand. *Food Chem.* 2021;334:127402. doi: [10.1016/j.foodchem.2020.127402](https://doi.org/10.1016/j.foodchem.2020.127402).
 125. Kumar M, Ramanathan AL, Tripathi R, Farswan S, Kumar D, Bhattacharya P. A study of trace element contamination using multivariate statistical techniques and health risk assessment in groundwater of Chhaprola Industrial Area, Gautam Buddha Nagar, Uttar Pradesh, India. *Chemosphere.* 2017;166:135-45. doi: [10.1016/j.chemosphere.2016.09.086](https://doi.org/10.1016/j.chemosphere.2016.09.086).
 126. Cheng Z, Chen LJ, Li HH, Lin JQ, Yang ZB, Yang YX, et al. Characteristics and health risk assessment of heavy metals exposure via household dust from urban area in Chengdu, China. *Sci Total Environ.* 2018;619-620:621-9. doi: [10.1016/j.scitotenv.2017.11.144](https://doi.org/10.1016/j.scitotenv.2017.11.144).
 127. Caggiano R, Sabia S, Speranza A. Trace elements and human health risks assessment of finer aerosol atmospheric particles (PM1). *Environ Sci Pollut Res Int.* 2019;26(36):36423-33. doi: [10.1007/s11356-019-06756-w](https://doi.org/10.1007/s11356-019-06756-w).
 128. Sanaei F, Amin MM, Pourjam Alavijeh Z, Azaripour Esfahani R, Sadeghi M, Shabankare Bandarrig N, et al. Health risk assessment of potentially toxic elements intake via food crops consumption: Monte Carlo simulation-based probabilistic and heavy metal pollution index. *Environ Sci Pollut Res Int.* 2021;28(2):1479-90. doi: [10.1007/s11356-020-10450-7](https://doi.org/10.1007/s11356-020-10450-7).
 129. Shalaby SE, Abdou GY, El-Metwally IM, Abou-Elella G. Health risk assessment of pesticide residues in vegetables collected from Dakahlia, Egypt. *J Plant Prot Res.* 2021;61(3):254-64. doi: [10.24425/jppr.2021.137951](https://doi.org/10.24425/jppr.2021.137951).
 130. Salehzadeh H, Maleki A, Rezaee R, Shahmoradi B, Ponnet K. The nitrate content of fresh and cooked vegetables and their health-related risks. *PLoS One.* 2020;15(1):e0227551. doi: [10.1371/journal.pone.0227551](https://doi.org/10.1371/journal.pone.0227551).
 131. Lemos J, Sampedro MC, de Ariño A, Ortiz A, Barrio RJ. Risk assessment of exposure to pesticides through dietary intake of vegetables typical of the Mediterranean diet in the Basque Country. *J Food Compos Anal.* 2016;49:35-41. doi: [10.1016/j.jfca.2016.03.006](https://doi.org/10.1016/j.jfca.2016.03.006).
 132. Maurice L, López F, Becerra S, Jamhoury H, Le Menach K, Dévier MH, et al. Drinking water quality in areas impacted by oil activities in Ecuador: associated health risks and social perception of human exposure. *Sci Total Environ.* 2019;690:1203-17. doi: [10.1016/j.scitotenv.2019.07.089](https://doi.org/10.1016/j.scitotenv.2019.07.089).
 133. Lozowicka B. Health risk for children and adults consuming apples with pesticide residue. *Sci Total Environ.* 2015;502:184-98. doi: [10.1016/j.scitotenv.2014.09.026](https://doi.org/10.1016/j.scitotenv.2014.09.026).
 134. Coscollà C, López A, Yahyaoui A, Colin P, Robin C, Poinsignon Q, et al. Human exposure and risk assessment to airborne pesticides in a rural French community. *Sci Total Environ.* 2017;584-585:856-68. doi: [10.1016/j.scitotenv.2017.01.132](https://doi.org/10.1016/j.scitotenv.2017.01.132).
 135. Nougadère A, Sirot V, Cravedi JP, Vasseur P, Feidt C, Fussell RJ, et al. Dietary exposure to pesticide residues and associated health risks in infants and young children - results of the French infant total diet study. *Environ Int.* 2020;137:105529. doi: [10.1016/j.envint.2020.105529](https://doi.org/10.1016/j.envint.2020.105529).