

Investigation of heavy metals accumulation in the soil and pine trees

Javad Kharkan^{ID}, Mohammad Hossein Sayadi^{ID}, Mohammad Reza Rezaei^{ID}

Department of Environmental Sciences, School of Natural Resources and Environment, University of Birjand, Birjand, Iran

Abstract

Background: Human activities related to workshops in the cities contribute to the release of heavy metals into the environment, which pose serious risks to the environment and to human health. The aim of the present study was to evaluate the concentration of lead (Pb), iron (Fe), and manganese (Mn) in the pine trees and soil in various land uses of Birjand city, Iran.

Methods: The sampling stations were randomly selected from different land uses including parks, streets, carwashes, car repair shops, and car smooth shops in Birjand city. The pine trees (skin and leaves) and soil samples were collected from 15 stations located at different and uses in 2017. To determine the concentration of heavy metals, atomic absorption spectrometer (Contr AA 700) was used.

Results: It was revealed that the mean concentration of Pb, Fe and Mn in residential soil was 1.79, 419.39, and 30.76 mg/kg, respectively. Moreover, the Pb, Fe, and Mn concentration in pine skin and leaves was 0.63 – 0.18, 23.05–9.84, and 10.05–3.13 mg/kg, respectively. The geo-accumulation index (I_{geo}) mean of the study areas demonstrated a descending trend for Fe (16.31 mg/kg) < Mn (8.86 mg/kg) < Pb (0.41 mg/kg). Pb showed the highest transfer factor in the parks and streets followed by Mn and Fe. In the car repair and smooth shops, the highest transfer factor pertained to Pb and the lowest one pertained to Fe. The statistical analysis indicated that there was a significant difference in the Fe and Mn concentrations among various land uses ($P < 0.05$).

Conclusion: According to the results, the soils of car repair and smooth shops as well as carwashes in Birjand are becoming polluted by Pb, Fe, and Mn. Although, it does not threaten the city ecosystem, but with passage of time, these measures will be accumulated due to the soil alkalinity and will reach critical levels.

Keywords: Lead, Iron, Manganese, Soil, Cities

Citation: Kharkan J, Sayadi MH, Rezaei MR. Investigation of heavy metals accumulation in the soil and pine trees. Environmental Health Engineering and Management Journal 2019; 6(1): 17–25. doi: 10.15171/EHEM.2019.03.

Article History:

Received: 20 October 2018

Accepted: 12 January 2019

ePublished: 28 January 2019

*Correspondence to:

Mohammad Hossein Sayadi

Email: Mh_sayadi@birjand.ac.ir

Introduction

Environmental pollution by heavy metals has become a matter of attention on local, regional, and global scales due to their toxicity effects (1-3). Essential metals such as iron (Fe) and manganese (Mn) play a role in metabolic activity and are required for growth, development, and maintenance of body, but high concentrations of these metals have adverse health effects on different organs such as liver, heart, endocrine glands, and neurotoxic effect on the brain (4,5). In contrast, the non-essential metals such as lead (Pb) has no role in metabolic activities of organisms but at low concentrations, it has the potential to affect toxicity on different tissues of living organisms (6,7). Pb exposure can cause different health problems such as anemia, bone, renal, reproductive, nervous system disorders, neurohematological damages, and cancers (8).

The urban soil is the most important part of an urban environment. Various studies have been conducted in many countries to investigate the soil contamination by heavy metals (Pb, Fe, and Mn) in urban soils, and it was revealed that heavy metals concentrations were affected by several factors such as traffic activities, parks, and various workshops (9,10). Human activities related to workshops in the cities contribute to the release of heavy metals into the environment, which pose serious risks to the environment and human health. Brake wear, and exhaust gas are the most important sources for releasing of Pb from vehicles (11). The levels of soil contamination and plant species are two main factors that affect the dynamics of heavy metals in plant-soil interactions (12). The roadside plants' leaves or stems may also absorb heavy metals through atmospheric deposit or road runoff (13,14).



Different parts of the plant contain different amounts of heavy metals, so that the highest concentrations of metals are found in roots, stems, and leaves (15).

Some trees such as pine have been used as biological indicators for monitoring heavy metals in the environmental pollution, because these trees are extensively dispersed in most cities as the main plant species with long lifetime (16). It is interesting to note that the dominant trees in the city of Birjand is the pine trees. The vehicles, carwashes, car smooth units, and car repair shops release heavy metals in the urban environment. Since, there has not been a study on the concentration of heavy metals in pine trees (skin and leaves) in the climatic conditions of the region and different land uses, therefore, this study aimed to determine the concentration of Pb, Fe, and Mn in the soil and pine trees at different land uses in Birjand city such as different parks, streets, carwashes, car repair shops, and car smooth shops.

Materials and Methods

Study area

Birjand city is the capital of South Khorasan province in the eastern part of Iran. Birjand city lies on the border with Afghanistan and it is about 1491 m from the sea level. At 2016 census, its population was 221 756 consisting 60 240 families. This city is located at a latitude of 32° 87' N and longitude of 59° 21' E with cold semi-arid climates (Figure 1). Moreover, Birjand city from the middle of April to December has a dry season with 171 mm rainfall and 255 sunny days in the year (17).

Sampling and laboratory analysis

The total of 15 soil and pine trees (skin and leaf) samples were randomly collected from different land uses in Birjand city such as car repair and smooth shops, carwashes, streets, and parks with three replicates and

transferred to the Central Laboratory of Environmental Pollutants of Birjand University. The pH of soil saturation extract was analyzed using a pH meter. Electrical conductivity (EC) of soil and soil texture was determined by an EC meter and hydrometer method, respectively (18). To measure pH, 5 g of each soil sample was placed into a 50 cc Erlenmeyer flask. Distilled water was added to the sample (1:25 dilution) and kept for 12 hours at ambient temperature. Then, pH was measured by the pH meter. Moreover, to determine the EC, 150 g of the screened soil (2 mm) was poured into a container and saturated with 150 mL distilled water (1:1). After saturation, its extract was obtained by Bacchini funnel. A filter paper was placed into the funnel and an extraction glass was kept under the funnel. The extracted soil was collected in the extraction glass. The physiochemical characteristics of the extracted soil samples were reported in Table 1.

Determination of the heavy metals concentration

To determine the concentration of heavy metals, 1 g of soaked and sieved soil was placed in an Erlenmeyer flask, then 16 cc acid [a mixture of 4 cc nitric acid (65%) and 12 cc hydrochloric acid (37%)] was added to the flask. For acid digestion, the flask was placed on the sand bath at 100°C for 6 to 7 hours. At the specified time period, 4 cc per chloric acid was added to the flask placed on the sand bath. After evaporation of 3 cc acid, the sample was flattened by a plastic funnel on the filter paper (18,19).

Determination of heavy metals concentration in pine trees

To determine heavy metals concentration in the plant texture, the samples were dried in an oven at 70°C for 48 hours, then, the samples were ground by a Chinese oven for better dissolving in the acid, finally, 1 g of the sample was placed into an Erlenmeyer flask for sample digestion. For chemical digestion of the samples, nitric acid (HNO_3)




Figure 1. Sampling site map in different parts of Birjand city in eastern Iran.

Table 1. Physiochemical characteristics of the studied soil samples

Land Use	pH Soil	Soil EC (MSCM-1)
Park	7.73	2.165
Street	7.43	6.67
Carwash	7.5	5.38
Car repair shop	8.5	4.98
Car smooth shop	8.2	6.76

65% and per chloric acid 70%-72% (5:1.5) were used, the samples were then placed on the sand bath at 100°C and heated until the color of the solution was clear. After digestion, the samples were placed at room temperature for cooling, then, the samples were diluted to 50 cc with double distilled water and flattened by filter paper (20). Finally, the samples were analyzed by graphite furnace atomic absorption spectroscopy (ContrAA 700) (with a stock concentration of 1 mg/mL and compute calibration adjusted $R^2 = 0.99$). The detection limits for each metal were as follows: Pb = 0.03, Mn = 0.10, and Fe = 0.11, and mean recovery of 98.4, 97.8, and 96%, respectively, were obtained.

Geo-accumulation index

Geo-accumulation index (I_{geo}) was used to evaluate soil pollution by comparing the current heavy metals concentration in the soil with their pre-industrial/background concentration. This index is used to determine the pollution degree and the amount of human factors affecting the soil and sediment (21) and is evaluated through Eq. (1):

$$I_{geo} = \log_2 \left[\frac{C_n}{1.5B_n} \right] \quad (1)$$

where C_n is the measured concentration of element in the sample and B_n is the concentration of the same element in the sample. To correct the effects of soil material through human activities, the correction coefficient of 1.5 was used. According to this index, the soils are categorized into 7 different groups as below (22).

$I_{geo} \leq 0$ (non-contaminated), $1 < I_{geo} < 0$ (non-contaminated to slightly infected), $2 < I_{geo} < 1$ (slightly infected), $3 < I_{geo} < 2$ (slightly infected to very polluted), $4 < I_{geo} < 3$ (very polluted), $5 < I_{geo} < 4$ (very polluted to severely infected), and $5 \leq I_{geo}$ (severely infected).

Contamination factor index

According to contamination factor (CF) index, the enrichment of metals in relation to the background concentrations can be evaluated and the degree of soil pollution can be determined (23). The contamination factor is determined using Eq. (2):

$$CF = [C]_{\text{sample}}/[C]_{\text{background}} \quad (2)$$

Where CF is contamination factor, (C) sample is the concentration of considered element, and (C) background is the concentration of basic element in the sample. In terms of contamination by heavy metals, soils are divided into 4 groups based on the contamination factor.

$1 > CF$ (lowly polluted), $3 < CF < 1$ (moderately polluted), $6 \leq CF < 3$ (highly polluted), and $6 \leq CF$ polluted (24,25). In this study, the sample was collected from the control soil (i.e. an area without any human activity) and from the depth of one and a half meters. The background concentration of Pb, Fe, and Mn was 0.26, 305.4, and 23.94, respectively.

Pollution load index

To evaluate the severity of contamination, the pollution load index (PLI) was used. Using PLI, the effects of total contamination by different metals is evaluated. The advantage of PLI over other indicators is that the pollution risk index of all the studied metals is determined in the area. According to this index, the standardized quality of soil is divided into 5 levels (26).

Highly polluted ($P \leq 0.7$), clean contaminated ($0.7 < P \leq 1$), slightly polluted ($1 < P \leq 2$), moderately polluted ($2 < P \leq 3$), and highly polluted ($P > 3$).

The pollution coefficient of Pb is evaluated based on the Eq. (3):

$$PLI = \sqrt[n]{CF_1 \times CF_2 \times CF_3 \times \dots \times CF_n} \quad (3)$$

Enrichment factor

The evaluation of the soils pollution was made by using enrichment factor to overcome the difficulties in making an overall assessment of the degree of metal contamination in soils. This factor is evaluated based on the Eq. (4)(27).

$$EF = \frac{\left[\frac{C_x}{C_{ref}} \right]_{\text{Sample}}}{\left[\frac{C_x}{C_{ref}} \right]_{\text{Background}}} \quad (4)$$

Where EF is enrichment factor, C_x is the concentration of measured element in the sample, and C_{ref} is the concentration of the reference element (Fe). The reference element is an element that has the geological origin. In the environmental studies, Sr, Al, Fe, Ti, and Zr are usually used as the reference elements (28). The degree of heavy metals pollution can be divided into 5 groups (29): $EF < 2$ (low pollution level), $2 \leq EF < 5$ (moderate pollution level), $5 \leq EF < 20$ (high pollution level), $20 \leq EF < 40$ (very high pollution level), and $40 \leq EF$ (severe pollution level).

Potential ecological risk

The index of potential ecological risk (Er) is used to evaluate the environmental potential risks of metals in the soil (30). In this study, the potential Er of heavy metals was determined using the following equations:

$$C_j^i = C^i / C_j^i Er = T_n^i \times C_j^i \quad (5)$$

Where C^i is the concentration of measured metal in studied soil sample, C_j^i is the reference of background value from this element, and T_n^i is the toxicity index of the heavy metal (Pb:5) (31). Er shows the potential ecological risk of Pb. The pollution level is divided into 5 levels based on the ecological risk. $Er < 40$ (low potential Er), $40 \leq Er < 80$ (moderate potential Er), $80 \leq Er < 160$ (significant potential Er), $160 \leq Er < 320$ (high potential Er), $320 \leq Er$ (very high potential Er).

Transfer factor of heavy metals

Transfer factor is described as the ratio of metal concentration in tree organ ($C_{part\ of\ tree}$) and concentration of the same metal in the soil (C_{soil}) (32). This factor is used to measure the capability of the tree to absorb the metal from the soil and is evaluated based on the Eq. (6):

$$TF = C_{Part\ of\ tree} / C_{soil} \quad (6)$$

Statistical analysis

Data were analyzed using SPSS version 20. The concentrations of heavy metals (Pb, Mn, and Fe) in soil and pine's skin and leaf were compared in different stations using one-way analysis of variance (ANOVA). Moreover, the Pearson correlation coefficient was used to assess the relationship among heavy metals in the soil of the study area. Data were expressed as mean \pm standard deviation (mean \pm SD).

Results

The solubility of heavy metals to dissolve and their access to plants depend on the physicochemical characteristics of the soil (e.g. acidity and texture) that are considered essential among the effective factors used in plants to absorb the heavy metals. Nevertheless, soil is usually considered as the most important factor (33). Mean pH of 7.87 represents an alkaline condition for all soil samples (Table 1). Effectually, with increasing pH, the access level to heavy metals decreases, so its concentration increases in the soil.

The average concentration of heavy metals Pb, Fe, and Mn was measured by atomic absorption spectrophotometer (AAS) in soil and pine trees (skin and leaf) in different land uses (Table 2). The average concentration of Pb, Fe, and Mn was 0.87, 150.79, and 14.65 mg/kg, respectively. The statistical analysis indicated that there are significant differences in Pb concentration in the soil and pine's skin among various land uses ($P > 0.05$). Also, there was a significant difference in the Fe concentration in pine's skin and leaf ($P < 0.01$). There were significant differences in Mn concentration in the soil and pine's skin ($P < 0.01$). The average concentration of Pb, Fe, and Mn in the soil was 1.79, 419.39, and 30.75 mg/kg, respectively. The average concentration of Pb, Fe, and Mn in the pine's skin and leaf was 0.63, 23.04, and 10.06 mg/kg and 0.17, 9.84, and 3.14 mg/kg, respectively. It is clear that, the concentration of these metals in different parts of pine exhibited a descending order of leaf < skin < soil.

The highest concentration of Pb belonged to car repair shop and the amount of Pb in different land uses was in a descending order of Park < Street < Car smooth shop < Carwash < Car repair shop. So, carwash and car repair shop significantly increased Pb in the urban environment. Pb had the highest concentration (4.56 mg/kg) in car repair shop and the lowest one in park (0.28). Fe and Mn in park demonstrated the highest concentration. Pb, Fe, and Mn showed the highest concentration in the soil and their concentration in pine's skin was more than the leaf. The statistical analysis showed that there are significant differences in Pb and Mn concentration in the soil and pine's skin among different land uses ($P < 0.01$). Also, there was a significant difference in the Fe concentration in pine's leaf among different land uses ($P < 0.01$).

The results of Pearson correlation coefficient showed that there was a strong correlation between the studied elements (Table 3). Strong correlation was observed between the elements that were contaminated from the common sources like wastes from car wash and repair or through soil materials. In a similar study, Bhuiyan et al evaluated the correlation of heavy metals by Pearson correlation coefficient and concluded that the metals that showed high correlation, probably had the same pollution

Table 2. The mean (\pm SD) of heavy metals (Mn, Fe, and Pb) in the soil and pine tree (skin and leaf) in different land uses (mg/kg)

Land use	Mn			Fe			Pb		
	Leaf	Skin	Soil	Leaf	Skin	Soil	Leaf	Skin	Soil
Park	2.44 \pm 1.66	23.81 \pm 3.91	42.77 \pm 3.14	15.54 \pm 1.16	53.2 \pm 7.99	478.8 \pm 73.83	0.15 \pm 0.40	0.45 \pm 0.08	0.28 \pm 0.07
Street	2.03 \pm 0.31	11.86 \pm 1.25	22.8 \pm 3.24	9.19 \pm 0.82	21.02 \pm 14.23	441.5 \pm 75.23	0.11 \pm 0.02	0.66 \pm 0.09	0.37 \pm 0.05
Carwash	2.81 \pm 0.54	4.26 \pm 0.30	27.21 \pm 1.28	9.16 \pm 0.92	15.83 \pm 1.89	480.9 \pm 51.16	0.14 \pm 0.06	0.18 \pm 0.06	2.05 \pm 0.43
Car repair shop	4.62 \pm 0.40	6.07 \pm 2.72	24.21 \pm 4.02	6.55 \pm 1.02	14.08 \pm 2.86	345.6 \pm 90.24	0.25 \pm 0.15	0.87 \pm 0.19	4.56 \pm 1.37
Car smooth shop	3.82 \pm 0.60	4.28 \pm 1.06	36.8 \pm 10.29	8.76 \pm 0.81	10.75 \pm 4.35	350.1 \pm 64.59	0.22 \pm 0.09	0.99 \pm 0.67	1.68 \pm 1.31
F value	4.57	7.80	40.43	11.94	3.07	1.28	2.59	15.32	36.89
P value	0.23	0.004	0.001	0.001	0.068	0.34	0.10	0.001	0.001

Table 3. The Pearson correlation among metal concentrations (Pb, Fe, and Mn) in soil

	Pb	Fe	Mn
Pb	1		
Fe	0.441**	1	
Mn	0.316*	0.832**	1

*At 95% level is significant; **At 99% level is significant.

sources (26).

Geo-accumulation index

According to Table 4, the mean of Igeo of the study areas exhibited an increasing order of Igeo Pb (0.41) < Igeo Mn (8.86) < Igeo Fe (16.31). As shown in this table, the Igeo value of both elements Fe and Mn was over 5, therefore, the area is very heavily polluted.

Contamination factor

The CF of Pb showed low (12.5%), moderate (31.25%), high (6.26%), and severe pollution (50%). Furthermore, Fe and Mn exhibited low (18.75%) and moderate pollution (81.25%), respectively (Table 5). As shown in Table 4, the CF mean in the study areas showed an increasing order of Mn (1.27) < Fe (1.35) < Pb (6.50).

Enrichment factor

The enrichment factor (EF) average for the studied heavy metals in the soil samples showed a descending order of Mn (1.27) < Fe (1.35) < Pb (6.51), therefore, the enrichment factor average for Pb was more than 1.5 (Table 6).

Potential ecological risk

In this study, the potential ecological risk (Er) for Pb was evaluated. The values of potential Er are presented in Table

Table 4. Descriptive statistical analyses of accumulation land index (Igeo) and contamination factor (CF) of the studied metals

Parameter	Mean	Maximum	Minimum	SD
Igeo Pb	0.41	4.68	0.08	1.65
Igeo Fe	16.31	16.77	15.82	0.32
Igeo Mn	8.86	9.48	8.31	0.4
CF Pb	6.5	23.42	0.87	6.73
CF Fe	1.35	1.8	0.93	0.29
CF Mn	1.27	1.86	0.83	0.38

Table 5. The contamination factor (CF) of heavy metals (Pb, Fe, and Mn) in the soil of different land uses

Metal	Very high pollution	High pollution	Moderate pollution	Low Pollution
Pb	50*	6.26	31.25	12.5
Fe	0	0	81.25	18.75
Mn	0	0	81.25	18.75

7. The results showed that the average potential Er for Pb was 34.36 and its low and significant values were 60% and 13.3%, respectively (Table 8). Thus, Pb is an effective factor in increasing the potential Er in the aforementioned and uses soil. This metal, which is found as a colored material and petroleum product and used in the car smooth and repair shop, is very dangerous.

Transfer factor

Transfer factor of the studied heavy metals in the soil and pine's skin and leaf is shown in Table 9. Pb showed the highest transfer factor in the parks and streets followed by Mn and Fe. In carwash, the transfer factor demonstrated a descending order of Fe < Pb < Mn. In the car repair and smooth shops, the highest transfer factor pertained to Pb and the lowest one to Fe. High value of transfer factor of Pb signifies high ability of pine to absorb Pb from the soil. In several plants, the concentration of Pb was hundred times more than the permissible limits of its accumulation in the plant.

Discussion

Different human activities like car workshops can cause soil pollution that leads to major environmental problems. After a long time, heavy metals can accumulate in the human body. The bivalent Pb is replaced in the body, instead of calcium ion, and this replacement permits Pb to remain in the body for a long time. Pb accumulation can disrupt the human circulatory system as well as kidneys. It also has serious negative effects on the physical development of children. The reasons for the use of Pb in car smooth, repair shop, and carwash is that it is one of the petroleum products and its derivatives are used not only in producing various colors, but also for disposal of motor oil from battery water (34). In a study in Kerman city, the most important factors that increase the heavy metals in the soil, especially Pb, Zn, and Cu, were traffic and its related activities such as gas stations, battery packs, oil changes, and car painting and smooth (35). The results of a study by Saboohi et al showed that the floor dust of Yazd battery factory workshops has high level of heavy metals concentration so that it has the potential environmental and health risks (36). The most dangerous

Table 6. The enrichment factor average for the heavy metals in the soil samples

Mn	Fe	Pb
1.27	1.35	6.51

Table 7. The results of statistical analysis of ecological risk (Er) for Pb in the studied stations

Ecological risk (Er)			
Mean	Maximum	Minimum	SD
34.36	117.2	4.33	34.07

Table 8. The results of statistical analysis of ecological risk (Er) for Pb in the studied stations

Very high ecological risk	High ecological risk	Significant ecological risk	Moderate ecological risk	Low ecological risk
0	0	2 (13.3%)	4 (26.6%)	9 (60%)

Table 9. Transfer factor of heavy metals (mean \pm SD) in different land uses

Land use	TF _{Mn}		TF _{Fe}		TF _{Pb}	
	Leaf	Skin	Leaf	Skin	Leaf	Skin
Park	0.6 \pm 0.4	0.56 \pm 0.6	0.33 \pm 0.05	0.11 \pm 0.01	0.56 \pm 0.7	1.68 \pm 0.1
Street	0.93 \pm 0.2	0.52 \pm 0.3	0.2 \pm 0.01	0.46 \pm 0.3	0.33 \pm 0.11	1.87 \pm 0.4
Carwash	0.10 \pm 0.2	0.16 \pm 0.05	0.2 \pm 0.01	0.3 \pm 0.01	0.07 \pm 0.2	0.9 \pm 0.1
Car repair shop	0.17 \pm 0.2	0.24 \pm 0.07	0.2 \pm 0.01	0.4 \pm 0.01	0.5 \pm 0.01	0.19 \pm 0.2
Car smooth shop	0.7 \pm 0.9	0.1 \pm 0.1	0.02 \pm 0.01	0.03 \pm 0.01	0.16 \pm 0.06	0.62 \pm 0.1

effects pertained to Fe, Zn, and Pb. The soil and trees near the parks and streets receive Pb pollution from vehicle gases. Kleckerova and Dočekalova reported that pollution of Pb and Ca in Berono's urban areas is due to human activities and urban activities like traffic (37). Yang et al in their study on urban soil found that Pb, Ca, and Cu had significant correlation so that the sources of pollution by these metals were traffic and industrial activities (30).

The close relationship among heavy metals like Pb, Fe, and Mn showing the same sources (38,39). The use of petroleum products and its derivatives such as thinner and use of different colors in car painting as well as disposal of motor oil from battery water are appropriate justifications for high Igeo values of heavy metals in car repair and smooth shops rather than parks and streets. Davashi and Azimzadeh in the study of heavy metals in the soil of refinery near Ghamishloo's shelter found that the use of petroleum products and its derivatives like gasoline are known as an appropriate justification for high value of Igeo_{Pb} in the soil around the refinery (40). Zhu et al in a study on Igeo index evaluated the degree of heavy metals pollution in the Nanjing's soil-China and categorized Pb, Zn, and Cu in the uninfected class, and besides, signified the effect of industrialization and urbanization on the degree of heavy metals pollution in the urban soils during the time (41). Therefore, the results of Igeo index showed that the pollution level in the study area was high.

Among aforementioned land uses, the workshop had the maximum share in Pb dispersion. The contamination factor assessment showed that the soil of workshops and smooth shops demonstrated high and severe pollution for Pb, and low and moderate pollution for Fe and Mn. Lu et al reported that the highest concentration of Pb and Mn was observed in the industrial areas and the lowest concentration of these elements was observed in the residential areas with low traffic (42). The enrichment factor average for Pb was more than 1.5, which signifies that the source of this metal in the soil was human activities (43). The contamination level of each element in the study areas based on the average of this factor revealed

that Pb had the highest pollution level, whereas Fe and Mn had the lowest one. In addition, the results showed that Pb enters the environment through human activities and it does not have a natural origin. Yang et al in their study on urban soils of Chung Chun found that Pb, Cu, Ca, and Hg have significant correlation and identified industrial activities and traffic as the sources of metals pollution (30). Taghipour et al reported that Pb, Zn, and Cu, respectively had the highest concentration in urban soil due to industrial activities (44). Therefore, the results are consistent with the results of this study wherein high level of Pb contamination was found in the soil of carwashes and repair workshops, so it is quite essential to decrease heavy metals levels to inhibit the potential hazards of these metals in the environment. Various studies of ecological risk assessment showed that Pb caused various diseases in humans like kidney problems, liver diseases, and cancer (45,46). Thus, Pb can affect the local ecological function in the long run and currently, it does not have the ability to create health and environmental problems (47-49). So, it is important to consider the levels of heavy metals to inhibit the potential hazards of these metals in the environment. The transfer factor shows that pine can be used to refine the polluted areas. Lebeau et al reported that the transfer factor of elements from terrestrial to aerial organs can be used to identify the over accumulated organs (50). Some researchers used the ratio of heavy metals concentration in the aerial part/root to describe the plant resistance and reaction against high levels of metals in the soil wherein this ratio is greater than one in plants containing more than one reagent, and less than one in plants having less than one reagent (51).

Conclusion

According to the results, the soils of car repair and smooth shops as well as carwashes in Birjand are becoming polluted by Pb, Fe, and Mn. Although, it does not threaten the city ecosystem, but with passage of time, these measures will be accumulated due to the soil alkalinity and will reach critical levels. The soil and plant surrounding the

streets and parks are contaminated by Pb from vehicles gases. Effectually, Pb concentration in the soil and plant increases with traffic volume and decreases with the distance from the main roads and streets. However, in this study, the contamination factors like accumulation land index and contamination factor were evaluated. The contamination load and enrichment factor showed that Pb showed moderate to high pollution, whereas Fe and Mn represented low to moderate pollution in the car repair and smooth shops. According to the ecological risk index, Pb, Fe, and Mn in the aforementioned land uses doesn't have the ability to create the health and environmental problems. However, to prevent these problems, the potential hazards of these metals in the environment should be considered. In parks and streets, the transfer factor more than 1 signifies the high ability of pine to absorb Pb from the soil. So, it can be safely concluded that pine can be used to refine the contaminated areas in parks and streets.

Acknowledgements

This article is a part of a Master's thesis sponsored by Birjand University. The authors would like to gratitude Birjand University for providing access to the laboratory facilities to perform this study.

Ethical issues

The authors certify that all data collected during the study are as stated in the manuscript, and no data from the study has been or will be published separately elsewhere.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors participated in the data collection, analysis, and interpretation. All authors critically reviewed, refined, and approved the manuscript.

References

1. Sayadi MH, Rezaei MR, Rezaei A. Fraction distribution and bioavailability of sediment heavy metals in the environment surrounding MSW landfill: a case study. *Environ Monit Assess* 2015; 187(1): 4110. doi: 10.1007/s10661-014-4110-1.
2. Maleki A, Azadi NA, Mansouri B, Majnoni F, Rezaei Z, Gharibi F. Health risk assessment of trace elements in two fish species of Sanandaj Gheshlagh Reservoir, Iran. *Toxicol Environ Health Sci* 2015; 7(1): 43-9. doi: 10.1007/s13530-015-0219-z.
3. Mansouri B, Salehi J, Etebari B, Moghaddam HK. Metal concentrations in the groundwater in Birjand flood plain, Iran. *Bull Environ Contam Toxicol* 2012; 89(1): 138-42. doi: 10.1007/s00128-012-0630-y.
4. Gentry PR, Van Landingham C, Fuller WG, Sulsky SI, Greene TB, Clewell HJ 3rd, et al. A tissue dose-based comparative exposure assessment of manganese using physiologically based pharmacokinetic modeling-The importance of homeostatic control for an essential metal. *Toxicol Appl Pharmacol* 2017; 322: 27-40. doi: 10.1016/j.taap.2017.02.015.
5. Rehman IU, Ishaq M, Ali L, Khan S, Ahmad I, Din IU, et al. Enrichment, spatial distribution of potential ecological and human health risk assessment via toxic metals in soil and surface water ingestion in the vicinity of Sewakht mines, district Chitral, Northern Pakistan. *Ecotoxicol Environ Saf* 2018; 154: 127-36. doi: 10.1016/j.ecoenv.2018.02.033.
6. Sayadi MH, Torabi S. Geochemistry of soil and human health: a review. *Pollut Res* 2009; 28(2): 257-62.
7. Li R, Tang C, Cao Y, Jiang T, Chen J. The distribution and partitioning of trace metals (Pb, Cd, Cu, and Zn) and metalloid (As) in the Beijiang River. *Environ Monit Assess* 2018; 190(7): 399. doi: 10.1007/s10661-018-6789-x.
8. Sayadi MH, Rezaei A, Sayyed MR. Grain size fraction of heavy metals in soil and their relationship with land use. *Proc Int Acad Ecol Environ Sci* 2017; 7(1): 1-11.
9. Saeedi M, Hosseinzadeh M, Jamshidi A, Pajooheshfar SP. Assessment of heavy metals contamination and leaching characteristics in highway side soils, Iran. *Environ Monit Assess* 2009; 151(1-4): 231-41. doi: 10.1007/s10661-008-0264-z.
10. Chen X, Xia X, Zhao Y, Zhang P. Heavy metal concentrations in roadside soils and correlation with urban traffic in Beijing, China. *J Hazard Mater* 2010; 181(1-3): 640-6. doi: 10.1016/j.jhazmat.2010.05.060.
11. Yan X, Zhang F, Zeng C, Zhang M, Devkota LP, Yao T. Relationship between heavy metal concentrations in soils and grasses of roadside farmland in Nepal. *Int J Environ Res Public Health* 2012; 9(9): 3209-26. doi: 10.3390/ijerph9093209.
12. Hasanvand H, Ghasemi Aghbash F, Solgi E, Pazhohan I. The distance from road effects on heavy metals accumulation in soil and leaves of Persian oak trees (*Quercus brantii*) in Aleshtar- Khorramabad highway. *Journal of Forest Research and Development* 2018; (4)1: 29-41. [In Persian].
13. Bi X, Ren L, Gong M, He Y, Wang L, Ma Z. Transfer of cadmium and lead from soil to mangoes in an uncontaminated area, Hainan Island, China. *Geoderma* 2010; 155(1-2): 115-20. doi: 10.1016/j.geoderma.2009.12.004.
14. Carrero JA, Goienaga N, Barrutia O, Artetxe U, Arana G, Hernandez A, et al. Diagnosing the impact of traffic on roadside soils through chemometric analysis on the concentrations of more than 60 metals measured by ICP/MS. In: Rauch S, Morrison GM, Monzón A, eds. *Highway and Urban Environment*. Dordrecht: Springer; 2010. p. 329-36. doi: 10.1007/978-90-481-3043-6_35.
15. Mirecki N, Agic R, Sunic L, Milenkovic L, Ilic ZS. Transfer factor as indicator of heavy metals content in plants. *Fresen Environ Bull* 2015; 24(11c): 4212-9.
16. Miri M, Ehrampoush MH, Ghaffari HR, Ebrahimi Aval H, Rezaei M, Najafpour F, et al. Atmospheric heavy metals biomonitoring using a local *Pinus eldarica* tree. *Health Scope* 2017; 6(1): e39241. doi: 10.17795/jhealthscope-39241.
17. Sayadi MH, Sayyed MR, Saptarshi PG. An assessment of the Chitgar river sediments for the short-term accumulation of the heavy metals (Cr, Co, Cd, Cu, Pb, Ni and Zn) from Tehran, Iran. *Pollut Res* 2008; 27(4): 627-34.
18. Sayadi MH, Rezaei MR, Rezaei A. Sediment toxicity and

- ecological risk of trace metals from streams surrounding a municipal solid waste landfill. *Bull Environ Contam Toxicol* 2015; 94(5): 559-63. doi: 10.1007/s00128-015-1518-4.
19. Huang SW, Jin JY. Status of heavy metals in agricultural soils as affected by different patterns of land use. *Environ Monit Assess* 2007; 139(1-3): 317. doi: 10.1007/s10661-007-9838-4.
 20. Whole Building Design Guide (WBDG). EPA 600/4-79-020 methods for chemical analysis of water and wastes. [cited 2018 Oct 2] Available from: <https://www.wbdg.org/ffc/epa/criteria/epa-600-4-79-020>.
 21. Shi G, Chen Z, Bi C, Li Y, Teng J, Wang L, et al. Comprehensive assessment of toxic metals in urban and suburban street deposited sediments (SDSs) in the biggest metropolitan area of China. *Environ Pollut* 2010; 158(3): 694-703. doi: 10.1016/j.envpol.2009.10.020.
 22. Odewande AA, Abimbola AF. Contamination indices and heavy metal concentrations in urban soil of Ibadan metropolis, southwestern Nigeria. *Environ Geochem Health* 2008; 30(3): 243-54. doi: 10.1007/s10653-007-9112-2.
 23. Naimi S, Ayoubi S. Vertical and horizontal distribution of magnetic susceptibility and metal contents in an industrial district of central Iran. *J Appl Geophy* 2013; 96: 55-66. doi: 10.1016/j.jappgeo.2013.06.012.
 24. Rashed MN. Monitoring of contaminated toxic and heavy metals, from mine tailings through age accumulation, in soil and some wild plants at Southeast Egypt. *J Hazard Mater* 2010; 178(1-3): 739-46. doi: 10.1016/j.jhazmat.2010.01.147.
 25. Islam S, Ahmed K, Al-Mamun H, Hoque F. Preliminary assessment of heavy metal contamination in surface sediments from a river in Bangladesh. *Environ Earth Sci* 2015; 73(4): 1837-48. doi: 10.1007/s12665-014-3538-5.
 26. Bhuiyan MA, Parvez L, Islam MA, Dampare SB, Suzuki S. Heavy metal pollution of coal mine-affected agricultural soils in the northern part of Bangladesh. *J Hazard Mater* 2010; 173(1-3): 384-92. doi: 10.1016/j.jhazmat.2009.08.085.
 27. Huang P, Li TG, Li AC, Yu XK, Hu NJ. Distribution, enrichment and sources of heavy metals in surface sediments of the North Yellow Sea. *Cont Shelf Res* 2014; 73: 1-13. doi: 10.1016/j.csr.2013.11.014.
 28. Abraham GM, Parker RJ. Assessment of heavy metal enrichment factors and the degree of contamination in marine sediments from Tamaki Estuary, Auckland, New Zealand. *Environ Monit Assess* 2008; 136(1-3): 227-38. doi: 10.1007/s10661-007-9678-2.
 29. Li X, Feng L. Multivariate and geostatistical analyses of metals in urban soil of Weinan industrial areas, Northwest of China. *Atmos Environ* 2012; 47: 58-65. doi: 10.1016/j.atmosenv.2011.11.041.
 30. Yang Z, Lu W, Long Y, Bao X, Yang Q. Assessment of heavy metals contamination in urban topsoil from Changchun City, China. *J Geochem Explor* 2011; 108(1): 27-38. doi: 10.1016/j.gexplo.2010.09.006.
 31. Lu A, Wang J, Qin X, Wang K, Han P, Zhang S. Multivariate and geostatistical analyses of the spatial distribution and origin of heavy metals in the agricultural soils in Shunyi, Beijing, China. *Sci Total Environ* 2012; 425: 66-74. doi: 10.1016/j.scitotenv.2012.03.003.
 32. Bose S, Jain A, Rai V, Ramanathan AL. Chemical fractionation and translocation of heavy metals in *Canna indica* L. grown on industrial waste amended soil. *J Hazard Mater* 2008; 160(1): 187-93. doi: 10.1016/j.jhazmat.2008.02.119.
 33. Erfanmanesh M, Afyuni M. Environmental Pollution Water, Soil and Air. Esfahan: Arkan; 2005. [In Persian].
 34. Kreith F, Tchobanoglous G. Handbook of Solid Waste Management. 2nd ed. New York: McGraw-Hill Education; 2002.
 35. HamzehMA, Mirzaee M, Mozaffari H. Concentrations of Pb, Zn, Cu and Cd and their distribution in the urban environment of Kerman, Iran. *Journal of Environmental Science and Technology* 2009; 11(3): 161-77. [In Persian].
 36. Sabouhi M, Nejadkoorki F, Azimzadeh HR, Taleshi A. Heavy metal pollution in the floor dust of yazd battery repairing workshops in 2016. *Iranian Journal of Health and Environment* 2016; 9(1): 127-38. [In Persian].
 37. Kleckeroová A, Dočekalová H. Dandelion plants as a biomonitor of urban area contamination by heavy metals. *Int J Environ Res* 2014; 8(1): 157-64. doi: 10.22059/ijer.2014.705.
 38. Li FL, Liu CQ, Yang YG, Bi XY, Liu TZ, Zhao ZQ. Natural and anthropogenic lead in soils and vegetables around Guiyang city, southwest China: a Pb isotopic approach. *Sci Total Environ* 2012; 431: 339-47. doi: 10.1016/j.scitotenv.2012.05.040.
 39. Dankoub Z, Ayoubi S, Khademi H, Lu SG. Spatial distribution of magnetic properties and selected heavy metals in calcareous soils as affected by land use in the Isfahan region, Central Iran. *Pedosphere* 2012; 22(1): 33-47. doi: 10.1016/S1002-0160(11)60189-6.
 40. Davashi L, Azimzadeh H. investigation on soil heavy metals pollution rate around Isfahan petroleum and refinery Industries near Ghameshlou wildlife refuge. *Journal of Natural Environment* 2015; 67(4): 403-14. [In Persian].
 41. Zhu J, Fang L, Nie Z, Gao X. Ecological risk assessment of heavy metal in urban area soil. 4th International Conference on Bioinformatics and Biomedical Engineering; 2010 Jun 18-20; Chengdu, China: IEEE; 2010. doi: 10.1109/ICBBE.2010.5516873.
 42. Lu X, Wang L, Lei K, Huang J, Zhai Y. Contamination assessment of copper, lead, zinc, manganese and nickel in street dust of Baoji, NW China. *J Hazard Mater* 2009; 161(2-3): 1058-62. doi: 10.1016/j.jhazmat.2008.04.052.
 43. Zhang J, Liu CL. Riverine composition and estuarine geochemistry of particulate metals in China--weathering features, anthropogenic impact and chemical fluxes. *Estuar Coast Shelf Sci* 2002; 54(6): 1051-70. doi: 10.1006/ecss.2001.0879.
 44. Taghipour M, Ayoubi S, Khademi H. Spatial variability of total Ni and Cu concentration in surface soils surrounding the Hamadan city using geostatistic technique. *Journal of Water and Soil Conservation* 2010; 17(2): 69-88. [In Persian].
 45. Hakanson L. An ecological risk index for aquatic pollution control. A sedimentological approach. *Water Res* 1980; 14(8): 975-1001. doi: 10.1016/0043-1354(80)90143-8.
 46. Bada BS, Olarinre TA. Characteristics of soils and heavy metal content of vegetation in oil spill impacted land in Nigeria. *Soilsproceedings* 2012; 1: 10-7.

47. Rashki Ghaleno O, Sayadi MH, Rezaei MR. Potential ecological risk assessment of heavy metals in sediments of water reservoir case study: Chah Nimeh of Sistan. *Proc Int Acad Ecol Environ Sci* 2015; 5(4): 89-96.
48. Ogunkunle CO, Fatoba PO. Pollution loads and the ecological risk assessment of soil heavy metals around a mega cement factory in southwest Nigeria. *Pol J Environ Stud* 2013; 22(2): 487-93.
49. Sayadi MH, Shabani M, Ahmadpour N. Pollution index and ecological risk of heavy metals in the surface soils of Amir-Abad area in Birjand city, Iran. *Health Scope* 2015; 4(1): e21137. doi: 10.17795/jhealthscope-21137.
50. Lebeau T, Braud A, Jezequel K. Performance of bioaugmentation-assisted phytoextraction applied to metal contaminated soils: a review. *Environ Pollut* 2008; 153(3): 497-522. doi: 10.1016/j.envpol.2007.09.015.
51. Zojaji F, Hassani AH, Sayadi MH. Bioaccumulation of chromium by *Zea mays* in wastewater-irrigated soil: an experimental study. *Proc Int Acad Ecol Environ Sci* 2014; 4(2): 62-7.