



Identification of clinically antibiotic resistant genes *Aac(3)-IIa* and *Aac(6')-Ib* in wastewater samples by multiplex PCR

Naser Samadi¹, Rahim Aali^{2*}, Esrafil Asgari³, Hamed Mirhosaeini⁴, Ali Shahriari⁵, Farhad Mahmoodi⁶, Fardin Nouri⁷, Asad Hamdi⁷, Farzad Fanaie⁷, Shahla Yosefi⁷, Saied Dadashi⁷

¹MSc of Cellular and Molecular Research Center, Faculty of medicine, Urmia University of Medical Sciences, Urmia, Iran

²Assistant Professor of Department of Environmental Health Engineering, Faculty of Nursing and Health, Urmia University of Medical Sciences, Urmia, Iran

³Instructor of Department of Environmental Health Engineering, Faculty of Nursing and Health, Urmia University of Medical Sciences, Urmia, Iran

⁴Instructor of Medical Microbiology, Department of Environmental Health Engineering, Faculty of Health, Arak University of Medical Sciences, Arak, Iran

⁵Assistant Professor of Department of Environmental Health Engineering, Environmental Health Research Center, Golestan University of Medical Sciences, Gorgan, Iran.

⁶PhD student of Department of Environmental Engineering, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran

⁷BSc of Environmental Health, Student Research Committee, Urmia University of Medical Sciences, Urmia, Iran

Abstract

Background: Aminoglycoside antibiotics are widely used in medical centers, particularly to treat infections. The resistance developed against these agents is a huge concern in health care. A number of researchers have reported that hospital and municipal wastewaters are among the most important dissemination sources of these agent into the environment. Some, however, do not agree with this opinion. In the present study, the prevalence of aminoglycoside resistance genes was investigated in raw and effluent wastewater from hospital and municipal wastewater treatment plants.

Methods: To conduct this descriptive-analytical study, 30 samples were taken according to sampling principles and cold cycle and transferred to the molecular laboratory. DNA was extracted by the freeze-thaw method using a kit (Promega). The genes *aac(3)-IIa* and *aac(6')-Ib* which code aminoglycoside resistance were examined in this study.

Results: The results indicated that the studied genes are present in 35% of urban and hospital wastewaters, and their frequency percentage is higher in hospital wastewater (52%) than urban wastewater (48%). The studied genes were identified in 61% of raw hospital wastewater samples; however, they were not detected in the output wastewater from the studied treatment plants.

Conclusion: Although, the studied genes were not detected in the final effluent, there is a high potential for their release into the environment. The current study demonstrated that the coding genes of aminoglycoside antibiotic resistance are present in raw urban and hospital wastewaters. In the case of improper exploitation of wastewater treatment plants, the output water can contaminate other environmental sections, such as soil and water resources, and result in the emission of these contaminants.

Keywords: Antibiotic resistance, Aminoglycosides, Urban and hospital wastewaters

Citation: Samadi N, Aali R, Asgari E, Mirhosaeini H, Shahriari A, Mahmoodi F, et al. Identification of clinically antibiotic resistant genes *Aac(3)-IIa* and *Aac(6')-Ib* in wastewater samples by multiplex PCR. Environmental Health Engineering and Management Journal 2015; 2(2): 47–52.

Article History:

Received: 14 January 2015

Accepted: 19 April 2015

ePublished: 20 June 2015

*Correspondence to:

Rahim Aali

Email: Aali@hlth.mui.ac.ir

Introduction

Excessive use of antibiotics in the treatment of humans and animals leads to antibiotic resistance (1-3). Antibiotic resistant bacteria (ARB) and antibiotic resistant genes (ARGs) cause serious problems in the treatment of infectious diseases. They enter the environment in different ways. Urban and hospital wastewaters have a high poten-

tial for the release of resistant bacteria and genes into the environment (4,5). Such release increases concerns about public health (4). Antibiotic resistance in the environment can be transferred to nonpathogenic bacteria from pathogenic bacteria and increase pressure on the aquatic ecology through physiological changes and population mobility (4-6). Research has shown that a large number of ARB and



ARGs, in particular hospital strains, have been found in regions where hospital wastewaters have been discharged (7). Importantly, some of these contaminated water resources are used for public water supplies. Some investigations have revealed that discharge of hospital wastewater is associated with the widespread dissemination of resistant bacteria in the environment (1). Other studies have demonstrated that urban wastewaters are important means through which resistant bacteria and genes are discharged into the environment (8). Some researchers have shown that resistant bacteria and genes are rarely removed from wastewater in treatment plant facilities (3,4). Some studies have even indicated that the number of resistant bacteria increases in such facilities (9).

Aminoglycosides are a group of medications that have similar chemical, antimicrobial, pharmacological, and toxic properties. Currently, this group includes streptomycin, gentamicin, tobramycin, kanamycin, amikacin, netilmicin, neomycin, sisomicin, and others (10). Aminoglycoside antibiotics are widely used in medical centers to treat life-threatening infections caused by gram-negative bacteria (10,11). Resistance to aminoglycosides is often caused by drug enzymatic changes by phosphorylation, adenylation, and acetylation of amine and hydroxyl groups, the most common mechanism of antibiotic resistance in bacteria (12). Among enzymes that change aminoglycosides, the *aac(6')-Ib* is predominant and causes resistance to many antibiotics. This gene was first identified in 1986 in *Klebsiella pneumoniae* isolates (13). Although the production of aminoglycoside-changing enzymes is the most common mechanism for producing resistance to aminoglycosides, the gram-negative bacteria that produce 16S rRNA methylase have become increasingly prevalent in recent years (14), and such widespread resistance to aminoglycosides has become a huge challenge to the medical profession (15). Filipova et al (16) and Chandrakanth et al (17) reported aminoglycoside resistant bacteria isolated from clinical environments. The *aac* gene is distributed in different microorganisms isolated from polluted waters (18-20). Szczepanowski et al (21) reported that aminoglycoside resistance genes prevailed in clinical compartments and can be spread to the environment through hospital wastewater. Considering the risk of these pollutant genes entering the environment, this study examined the prevalence of the *aac(3)-IIa* and *aac(6')-Ib* resistant genes in raw and wastewater effluent and determined the role of urban wastewater treatment plants in

removing/reducing *aac* genes.

Methods

Preparation of samples

A certain volume of the main samples (50 ml) was taken and centrifuged for 10 minutes at 6000 rpm. The sediment was extracted, 300 µl of distilled water was added, and the solution was subjected to freeze and thaw using liquid nitrogen and boiling water, each for 1 minute periodically at 4-5 stages (22).

DNA extraction

DNA was extracted using a Promega Wizard Genomic DNA purification kit, (Madison, WI) in 3 stages. In the lysing stage of the cell, pipetting was performed well after the addition of cell lyse solution (600 µl). The solution was then incubated at 80°C for 5 minutes. Finally, RNase was added, and the solution was incubated again for 15-60 minutes at 37°C. At the protein precipitation stage, 200 µl of precipitation solution was added and incubated for 5 minutes on ice, followed by centrifugation at 13000-16000 rpm for 3 minutes. At the DNA precipitation stage, the top liquid was transferred to a tube containing 600 µl of isopropanol at room temperature and mixed. Afterwards, 600 µl of 70% ethanol was added at room temperature and the mixture was centrifuged for 2 minutes at 13000-16000 rpm. The ethanol was then removed and the tubes dried under ambient conditions. DNA was suspended in 100 µl of distilled water and kept for 1 night at 4°C; it was ultimately maintained at -20°C until molecular experiments were performed. The concentration of the extracted DNA was read by spectrophotometry (Nanodrop ND 1000).

PCR method

The pair of primers used to amplify the *aac(3)-IIa* and *aac(6')-Ib* genes are listed in Table 1. PCR amplification was conducted at a volume of 25 µl containing 2.5 µl buffer 10X PCR, 1 µl of each primer, 0.75 µl of MgCl₂, 0.5 µl of dNTP, 0.35 µl of tag polymerase, 2 µl of the DNA template, and 14.9 µl of deionized water. All PCR experiments had positive and negative controls. The process was performed using the initial denaturation stage for 5 minutes at 94°C, 35 cycles at 94°C for 45 seconds (denaturation), at 53°C for 30 seconds (annealing), and at 72°C for 1 minute (extension). The final extension was performed at 72°C for 10 minutes. PCR products were investigated using electrophoresis (1.5%) and DNA safe stain. They were then de-

Table 1. Characteristics of primers

Primers	Target gene	Sequence (5'-3')	Amplified size (bp)	Annealing temperature (°C)	References
F R	<i>Aac(3)-IIa</i>	F: 5-CGGAAGGCAATAACGGAG-3 R: 5-TCGAACAGGTAGCACTGAG-3	749 bp	53°C	(23)
F R	<i>Aac(6')-Ib</i>	F: 5'-ATG ACT GAG CAT GAC CTT G-3' R: 5'-AAGGGTTAGGCAACTG-3'	524 bp	53°C	(23)

tected on Transluminar UV.

Data analysis

To compare the presence and absence of genes among various sites, the chi-square test was employed. McNemar test was utilized before and after each treatment plant.

Results

Amplified multiplex genes are illustrated in Figure 1. The results indicated that *aac(3)-IIa* and *aac(6')-Ib* genes were found in 35% of the samples (Figure 2A). The *aac(3)-IIa* (37%) was found in a larger number of samples than the *aac(6')-Ib* gene (33%) (Figure 2B). Distinctive analysis of urban and hospital wastewaters indicated that the detection percentage of the studied genes was higher in hospital wastewaters (52%) than in urban wastewaters (48%) (Figure 2C).

The results revealed that the *aac(3)-IIa* gene was present in 50% of the hospital wastewater samples, while *aac(6')-Ib* was observed in 42% of the samples. The studied genes were found in 28% of the urban wastewater samples. In-

vestigations of the input and output of urban wastewaters indicated that, generally, 56% of the studied genes were found in the input. *aac(3)-IIa* and *aac(6')-Ib* were presented in the raw samples 56% and 44%, respectively. The studied genes were found in none of the effluent of urban wastewater treatment plants, but in 61% of the wastewater samples from hospitals. One of the studied hospitals had its own wastewater treatment plant; the results demonstrated that 83% of samples from this plant's raw wastewater contained the studied genes, but none of the genes were found in the effluent.

Discussion

Aminoglycoside antibiotics are widely used in medical clinics, especially to treat life-threatening infections caused by gram-negative bacteria (24-26). Because of the widespread incidence of aminoglycoside resistance and the resulting complications for the medical field (24), the present study investigated two aminoglycoside-resistance genes. Clinical genes resistant to aminoglycosides were found to be present in municipal wastewaters. In their studies, Guardabassi et al (27), Heuer et al (18), and Aali et al (8) also indicated that the genes resistant to aminoglycosides exist in environmental resources. The results of this study revealed the lowest percentage of clinically genes. The results of other studies also confirm these findings. Jakobsen et al (28) found *aac* genes in wastewater isolates and showed a possible dissemination of *aac* from hospital to wastewater. Aali et al (8) reported an association between the lowest number of bacteria and aminoglycoside-resistance genes. The reason can be attributed to less use of aminoglycoside antibiotics. Recently, the consumption of some members of the aminoglycoside family (such as




Figure 1. Aminoglycoside resistance genes, *aac(3)-IIa* (749) bp and *aac(6')-Ib*(524 bp).




Figure 2. (A) Overall detection of ARGs, (B) total detection of *aac(3)-IIa* and *aac(6')-Ib*, and (C) differences between municipal and hospital wastewaters.

gentamycin) has been limited because of side effects; however, studies have indicated that resistance can develop even in low antibiotic concentrations (29). Heuer et al (18) attributed the diversity of the genes that code resistance to gentamycin (including *Aac3-I*) in different environments to the increased frequency of genetic transfer elements. The primary mechanism involved in resistance to gentamycin has been reported as enzyme deactivation, and over 50 types of enzyme have been identified. These enzymes, based on their biochemical action, include acetyltransferases, phosphotransferases, and nucleotidyl transferases (adenyl transferases) (16,17,30). These enzymes are coded by various genes, including *Aac*, *Aph*, and *Ant* (*Aad*). The mentioned enzymes and the coder genes have been identified in a wide range of isolated bacteria (*Aeromonas*, *Escherichia*, *Vibrio*, *Salmonella*, and *Listeria*) from hospital environments and contaminated waters (18-20). The *Aac3* gene has been identified as the coder of aminoglycoside-N-acetyl transferases in the microbial populations isolated from wastewater treatment facilities, river water, and agricultural regions (31,32). This can also be applicable to this study, yet complementary research is necessary to investigate the role of key factors in genetic transfer in the studied wastewaters.

The findings of the current study revealed that the percentages of *aac(3)-IIa* and *aac(6')-Ib* genes are higher in hospital wastewaters than in urban wastewaters. These results are consistent with those of other researchers. The high rate of aminoglycoside-resistance genes in hospitals can be caused by common unnecessary prescriptions for patients. This also demonstrates the aminoglycoside resistant mobile genetic agents' movements between clinical and environmental bacteria (18). Plasmid and integrons have main roles in the dissemination of these agents (30). In their study of the *Aac3-1* gene, Aali et al (8) reported that the genes and bacteria resistant to aminoglycosides are more prevalent in hospital than urban wastewaters. Their low percentage in urban wastewaters can be related to process design and operation. The *aac(3)-IIa* gene was identified in hospital wastewaters more than the *aac(6')-Ib* gene. This can be due to bacteria carrying or better expression of the *aac(3)-IIa* gene.

In the current study, contrary to other studies, neither *aac(3)-IIa* nor *aac(6')-Ib* genes were detected in effluent wastewater samples from different wastewater treatment plants. Szczepanowski et al (21) identified many aminoglycoside resistance genes (eg, *aacA*, *aacA1*, *aacA4*, *aacA7*, *aacA29b*, *aacC1*, *aacC2*, *aacC3*, *aacC4*, *aac(3)-Id*, and *aac[69]-Im*) in the final effluent of wastewater treatment plants. The absence of the studied resistance genes in the effluent wastewater can be due to the effect of the plant's disinfection unit, the shock caused by a huge organic load, or the specific conditions of the wastewater treatment plants.

Conclusion

Human contact with resistant bacteria and genes has been a health concern associated with the management of diseases. This contact can occur through various means such as environmental resources. Hospital and urban wastewater plays a great role in the spread of resistant genes and bacteria. Wastewater treatment plants are very important as a controlling barrier against these contaminants entering environmental resources. Therefore, the design, operation, and continual monitoring of the effluent from treatment plants together with the monitoring of design parameters can be essential in the success of wastewater treatment plants in controlling these pollutants. Various parameters of treatment together with gene expression and the way genes are affected in the wastewater treatment process require further broad and extensive research. The presence of *aac(3)-IIa* and *aac(6')-Ib* genes in the wastewater samples of treatment plants indicates that conventional treatment approaches cannot guarantee the absence of these genes; thus it is suggested that more advanced methods be applied to control antibiotic-resistant genes.

Acknowledgments

The authors would like to thank the staff of the Cellular and Molecular Laboratory (Urmia Medicine Faculty) and Dr. Nikaeen, Head of the Advanced Environmental Molecular Laboratory (Department of Environmental Health Engineering), Isfahan University of Medical Sciences.

Ethical issues

The authors certify that all data collected during the study is presented in this manuscript, and no data from the study has been or will be published separately.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RA designed the study, NS (PCR setup), RA (DNA extraction and PCR setup), FF (testing), FN (testing), AH (testing), SD (testing) and SHY (testing) did the lab work, and ASH, HM, and FM performed the literature search and wrote the manuscript. All authors critically reviewed, refined, and approved the manuscript.

References

1. Kümmerer K. The presence of pharmaceuticals in the environment due to human use--present knowledge and future challenges. *J Environ Manageme* 2009; 90(8): 2354-66.
2. Tao R, Ying GG, Su HC, Zhou HW, Sidhu JP. Detection of antibiotic resistance and tetracycline resistance genes in Enterobacteriaceae isolated from the Pearl rivers in South China. *Environ Pollut* 2010;

- 158(6): 2101-9.
3. Kim S, Aga DS. Potential ecological and human health impacts of antibiotics and antibiotic-resistant bacteria from wastewater treatment plants. *J Toxicol Environ Health B Crit Rev* 2007; 10(8): 559-73.
 4. Munir M, Wong K, Xagorarakis I. Release of antibiotic resistant bacteria and genes in the effluent and biosolids of five wastewater utilities in Michigan. *Water Res* 2011; 45(2): 681-93.
 5. Gao P, Munir M, Xagorarakis I. Correlation of tetracycline and sulfonamide antibiotics with corresponding resistance genes and resistant bacteria in a conventional municipal wastewater treatment plant. *Sci Total Environ* 2012; 421-422: 173-83.
 6. Pruden A, Larsson DG, Amézquita A, Collignon P, Brandt KK, Graham DW, et al. Management Options for Reducing the Release of Antibiotics and Antibiotic Resistance Genes to the Environment. *Environ Health Perspect* 2013; 121(8): 875.
 7. Rahube TO, Yost CK. Antibiotic resistance plasmids in wastewater treatment plants and their possible dissemination into the environment. *Afr J Biotechnol* 2010; 9(54): 9183-90.
 8. Aali R, Nikaeen M, Khanahmad H, Hassanzadeh A. Monitoring and comparison of antibiotic resistant bacteria and their resistance genes in municipal and hospital wastewaters. *Int J Prev Med* 2014; 5(7): 887-94.
 9. Aali R, Nikaeen M, Khanahmad H, Hejazi Z, Kazemi M, Hassanzadeh A. Occurrence of tetracycline resistant bacteria and resistance gene(*tetW*) in hospital and municipal wastewaters. *Fresenius Environmental Bulletin* 2014; 23(10a): 2560-6.
 10. Klare I, Konstabel C, Badstübner D, Werner G, Witte W. Occurrence and spread of antibiotic resistances in *Enterococcus faecium*. *Int J Food Microbiol* 2003; 88(2-3): 269-90.
 11. Bradford PA. Extended-spectrum beta-lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clin Microbiol Rev* 2001; 14(4): 933-51.
 12. Wright GD. Antibiotic resistance in the environment: a link to the clinic? *Curr Opin Microbiol* 2010; 13(5): 589-94.
 13. Savic M, Lovric J, Ilic Tomic T, Vasiljevic B, Conn LG. Determination of the target nucleosides for members of two families of 16S rRNA methyltransferases that confer resistance to partially overlapping groups of aminoglycoside antibiotics. *Nucleic Acids Res* 2009; 37(16): 5420-31.
 14. Yamane K, Wachino JI, Suzuki S, Shibata N, Kato H, Shibayama K, et al. 16S rRNA Methylase producing, Gram-negative Pathogens. *Emerg Infect Dis* 2007; 13(4): 642-6.
 15. Jawetz E, Melnick JL, Adelberg EA. Review of Medical Microbiology. 22nd ed. Los Altos: Appleton & Lange; 2007.
 16. Filipova M, Bujdakova H, Drahovska H, Lišková A, Hanzen J. Occurrence of aminoglycoside-modifying-enzyme genes *aac* (6')-aph (2''), *aph* (3'), *ant* (4') and *andant* (6) in clinical isolates of *Enterococcus faecalis* resistant to high-level of gentamicin and amikacin. *Folia Microbiol (Praha)* 2006; 51(1): 57-61.
 17. Chandrakanth RK, Raju S, Patil S. Aminoglycoside-resistance mechanisms in multidrug-resistant *Staphylococcus aureus* clinical isolates. *Curr Microbiol* 2008; 56(6): 558-62.
 18. Heuer H, Krögerrecklenfort E, Wellington EM, Egan S, van Elsas JD, van Overbeek L, et al. Gentamicin resistance genes in environmental bacteria: prevalence and transfer. *FEMS Microbiol Ecol* 2002; 42(2): 289-302.
 19. Tennstedt T, Szczepanowski R, Braun S, Pühler A, Schlüter A. Occurrence of integron-associated resistance gene cassettes located on antibiotic resistance plasmids isolated from a wastewater treatment plant. *FEMS Microbiol Ecol* 2003; 45(3): 239-52.
 20. Tennstedt T, Szczepanowski R, Krahn I, Pühler A, Schlüter A. Sequence of the 68,869 bp IncP-1[alpha] plasmid pTB11 from a waste-water treatment plant reveals a highly conserved backbone, a Tn402-like integron and other transposable elements. *Plasmid* 2005; 53(3): 218-38.
 21. Szczepanowski R, Linke B, Krahn I, Gartemann KH, Gützkow T, Eichler W, et al. Detection of 140 clinically relevant antibiotic resistance genes in the plasmid metagenome of wastewater treatment plant bacteria showing reduced susceptibility to selected antibiotics. *Microbiology* 2009; 155: 2306-19.
 22. Asghari FB, Nikaeen M, Mirhendi H. Rapid monitoring of *Pseudomonas aeruginosa* in hospital water systems: a key priority in prevention of nosocomial infection. *FEMS Microbiol Lett* 2013; 343(1): 77-81.
 23. Samadi N, Hosainzadegan H, Akbari Qomi M. Assessment of *aac*(3')-IIa and *aac*(6')-Ib genes incidence involved in aminoglycoside resistance in *Klebsiella pneumoniae* isolated from Clinical samples in Urmia hospitals. *Jundishapur Journal of Microbiology* 2014: in press.
 24. Yu FY, Yao D, Pan JY, Chen C, Qin ZQ, Parsons C, et al. High prevalence of plasmid-mediated 16S rRNA methylase gene *rmtB* among *Escherichia coli* clinical isolates from a Chinese teaching hospital. *BMC Infect Dis* 2010; 10: 184.
 25. Bogaerts P, Galimand M, Bauraing C, Deplano A, Vanhoof R, De Mendonca R, et al. Emergence

- of ArmA and RmtB aminoglycoside resistance 16S rRNA methylases in Belgium. *J Antimicrob Chemother* 2007; 59(3): 459-64.
26. Zhang Y, Zhou H, Shen XQ, Shen P, Yu YS, Li LJ. Plasmid-borne armA methylase gene, together with blaCTX-M-15 and blaTEM-1 in a *Klebsiella oxytoca* isolate from China. *J Med Microbiol* 2008; 57(Pt 10): 1273-6.
 27. Guardabassi L, Lo Fo Wong DM, Dalsgaard A. The effects of tertiary wastewater treatment on the prevalence of antimicrobial resistant bacteria. *Water Res* 2002; 36(8): 1955-64.
 28. Jakobsen L, Sandvang D, Hansen LH, Bagger-Skjøt L, Westh H, Jørgensen C, et al. Characterisation, dissemination and persistence of gentamicin resistant *Escherichia coli* from a Danish university hospital to the waste water environment. *Environ Int* 2008; 34(1): 108-15.
 29. Bouki C, Venieri D, Diamadopoulos E. Detection and fate of antibiotic resistant bacteria in wastewater treatment plants: a review. *Ecotoxicol Environ Saf* 2013; 91: 1-9.
 30. Zhang XX, Zhang T, Fang H. Antibiotic resistance genes in water environment. *Appl Microbiol Biotechnol* 2009; 82(3): 397-414.
 31. Park J, Lee J, Oh J, Jeong Y, Cho J, Joo H, et al. Antibiotic selective pressure for the maintenance of antibiotic resistant genes in coliform bacteria isolated from the aquatic environment. *Water Sci Technol* 2003; 47(3): 249-53.
 32. Szczepanowski R, Krahn I, Linke B, Goesmann A, Pühler A, Schlüter A. Antibiotic multiresistance plasmid pRSB101 isolated from a wastewater treatment plant is related to plasmids residing in phytopathogenic bacteria and carries eight different resistance determinants including a multidrug transport system. *Microbiology* 2004; 150(11): 3613-30.